

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 April 2004 (29.04.2004)

PCT

(10) International Publication Number
WO 2004/035548 A1

(51) International Patent Classification⁷: C07D 235/06,
235/12, 401/06, 401/14, 403/06, 405/06, 405/14, A61K
31/4184, 31/4439, A61P 25/04, 25/16, 35/00

AstraZeneca R & D Montreal, 7171 Frederick-Banting, St.
Laurent, Montreal, Québec H4S 1Z9 (CA).

(21) International Application Number:
PCT/SE2003/001604

(74) Agent: GLOBAL INTELLECTUAL PROPERTY; As-
traZeneca AB, S-151 85 Södertälje (SE).

(22) International Filing Date: 15 October 2003 (15.10.2003)

(25) Filing Language: English

(26) Publication Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,
MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT,
RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(30) Priority Data:
0203070-8 16 October 2002 (16.10.2002) SE

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): AS-
TRAZENECA AB [SE/SE]; S-151 85 Södertälje (SE).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): PAGÉ, Daniel
[CA/CA]; AstraZeneca R & D Montreal, 7171 Freder-
ick-Banting, St. Laurent, Montreal, Québec H4S 1Z9
(CA). WALPOLE, Christopher [GB/CA]; AstraZeneca
R & D Montreal, 7171 Frederick-Banting, St. Laurent,
Montreal, Québec H4S 1Z9 (CA). YANG, Hua [CA/CA];

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: NOVEL COMPOUNDS



(57) Abstract: Compounds of formula (I) wherein R^{F1} , R^{F2} , Z, R^1 , R^2 , Ar, X and Y are as defined in the specification as well as salts and pharmaceutical compositions including the compounds were prepared. These compounds are useful in therapy, in particular in the management of pain.

NOVEL COMPOUNDS

BACKGROUND OF THE INVENTION

1. Field of the invention

5 The present invention relates to compounds that are therapeutic, and more particularly, to compounds that are effective in treating pain, cancer, multiple sclerosis, Parkinson's disease, Huntington's chorea and/or Alzheimer's disease.

2. Discussion of Relevant Technology


10 Pain management has been studied for many years. Cannabinoid receptor (e.g., CB₁ receptor, CB₂ receptor) ligands such as agonists, antagonists and inverse agonists may produce relief of pain in a variety of animal models by interacting with CB₁ and/or CB₂ receptors. Generally, CB₁ receptors are located predominately in the central nervous system, whereas CB₂ receptors are located primarily in the periphery
15 and are primarily restricted to the cells and tissues derived from the immune system.

 While CB₁ receptor agonists and mixed (CB₁/CB₂) receptor agonists, such as tetrahydrocannabinol (THC) and opiate drugs, are effective in anti-nociception models in animals, they tend to exert many undesired CNS side-effects, e.g., psychoactive side effects and the abuse potential of opiate drugs. These undesired
20 CNS side effects are known to be mediated by the CB₁ receptors. In contrast, CB₂ receptor agonists may manage pain in humans or animals without causing those undesired CNS side effects due to the general location of CB₂ receptors and other factors.

 Therefore, there is a need for CB₂ receptor ligands such as agonists useful in
25 managing pain and/or treating other symptoms or diseases.

SUMMARY OF THE INVENTION

 In one aspect, the invention provides a compound of formula (I) or pharmaceutically acceptable salts thereof:



wherein

R^{F1} and R^{F2} are independently electron-withdrawing groups;

Z is selected from O= and S=;

R^1 is selected from C_{1-10} alkyl; C_{1-10} alkyl substituted by at least one of
 5 halogen, cyano, acetoxymethyl and nitro; C_{2-10} alkenyl; C_{2-10} alkenyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; C_{2-10} alkynyl; C_{2-10} alkynyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; R^3R^4N - C_{1-6} alkyl; $R^3R^4NC(=O)$ - C_{1-6} alkyl; R^3O - C_{1-6} alkyl; $R^3OC(=O)$ - C_{1-6} alkyl; $R^3C(=O)$ - C_{1-6} alkyl; $R^3C(=O)NR^3$ - C_{1-6} alkyl; $R^3R^4NSO_2$ - C_{1-6} alkyl; $R^3CSO_2N(R^4)$ - C_{1-6} alkyl;
 10 $R^3R^4NC(=O)N(R^5)$ - C_{1-6} alkyl; $R^3R^4NSO_2N(R^5)$ - C_{1-6} alkyl; aryl- C_{1-6} alkyl; aryl- $C(=O)$ - C_{1-6} alkyl; heterocyclyl- C_{1-6} alkyl; heterocyclyl- $C(=O)$ - C_{1-6} alkyl; substituted aryl- C_{1-6} alkyl; substituted aryl- $C(=O)$ - C_{1-6} alkyl; substituted heterocyclyl- C_{1-6} alkyl; substituted heterocyclyl- $C(=O)$ - C_{1-6} alkyl; and C_{1-10} hydrocarbylamino;

R^2 is selected from C_{1-6} alkyl, substituted C_{1-6} alkyl, C_{2-6} alkenyl, substituted
 15 C_{2-6} alkenyl, C_{2-6} alkynyl, substituted C_{2-6} alkynyl, C_{3-6} cycloalkyl, substituted C_{3-6} cycloalkyl, aryl, substituted aryl, and C_{5-6} heteroaryl, and substituted C_{5-6} heteroaryl;

R^3 , R^4 and R^5 are independently selected from -H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and a divalent C_{1-6} group that together with another divalent C_{1-6} group
 20 forms a portion of a ring;

X is a C_{1-10} divalent group that separates groups connected thereto by one or two atoms;

Ar is a C_{4-12} divalent aromatic group; and

Y is selected from -CH= and -N=.

25 Another aspect of the invention is the use of the compound of the invention as a medicament.


A further aspect of the invention is to use the compound of the invention in the treatment of pain, cancers, multiple sclerosis, Parkinson's disease, Huntington's chorea and/or Alzheimer's disease.

30

In a further aspect, the present invention provides a pharmaceutical composition comprising a compound of the invention and a pharmaceutically acceptable carrier.

In a further aspect, the present invention provides a method for the therapy of pain in a warm-blooded animal, comprising: administering to said animal in need of such therapy a therapeutically effective amount of a compound of the invention.

In a further aspect, the present invention provides a method of producing a compound comprising: reacting a compound represented by formula (II) with $R^2OArXCOA$:



wherein

R^{F1} and R^{F2} are independently electron-withdrawing groups;

Z is selected from O= and S=;

R^1 is selected from C_{1-10} alkyl; C_{1-10} alkyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; C_{2-10} alkenyl; C_{2-10} alkenyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; C_{2-10} alkynyl; C_{2-10} alkynyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; $R^3R^4N-C_{1-6}$ alkyl; $R^3R^4NC(=O)-C_{1-6}$ alkyl; R^3O-C_{1-6} alkyl; $R^3OC(=O)-C_{1-6}$ alkyl; $R^3C(=O)-C_{1-6}$ alkyl; $R^3C(=O)NR^3-C_{1-6}$ alkyl; $R^3R^4NSO_2-C_{1-6}$ alkyl; $R^3CSO_2N(R^4)-C_{1-6}$ alkyl; $R^3R^4NC(=O)N(R^5)-C_{1-6}$ alkyl; $R^3R^4NSO_2N(R^5)-C_{1-6}$ alkyl; aryl- C_{1-6} alkyl; aryl- $C(=O)-C_{1-6}$ alkyl; heterocyclyl- C_{1-6} alkyl; heterocyclyl- $C(=O)-C_{1-6}$ alkyl; substituted aryl- C_{1-6} alkyl; substituted aryl- $C(=O)-C_{1-6}$ alkyl; substituted heterocyclyl- C_{1-6} alkyl; substituted heterocyclyl- $C(=O)-C_{1-6}$ alkyl; and C_{1-10} hydrocarbylamino;

R^2 is selected from C_{1-6} alkyl, substituted C_{1-6} alkyl, C_{2-6} alkenyl, substituted C_{2-6} alkenyl, C_{2-6} alkynyl, substituted C_{2-6} alkynyl, C_{3-6} cycloalkyl, substituted C_{3-6} cycloalkyl, aryl, substituted aryl, and C_{5-6} heteroaryl, and substituted C_{5-6} heteroaryl;

R³, R⁴ and R⁵ are independently selected from -H, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, and a divalent C₁₋₆group that together with another divalent C₁₋₆group forms a portion of a ring;


X is a C₁₋₁₀divalent group that separates groups connected thereto by one or
5 two atoms;

A is selected from $-\text{OH}$, $-\text{Cl}$, $-\text{Br}$, and $-\text{I}$;

Ar is a C₄₋₁₂ divalent aromatic group; and

Y is selected from $-\text{CH}=\text{}$ and $-\text{N}=\text{}$.

10 In another aspect, the present invention provides a method of producing a compound comprising the step of reacting a compound represented by formula (III) with formaldehyde:



wherein

15 r and s are selected from 0, 1 and 2;

R¹⁰ is selected from C₁₋₆alkylene, -O-, and -NR¹¹-, wherein R¹¹ is a C₁₋₆alkyl;

R^{F1} and R^{F2} are independently electron-withdrawing groups;

X is a C₁₋₁₀divalent group that separates groups connected thereto by one or two atoms;

20 Ar is a C₄₋₁₂divalent aromatic group;

R² is selected from C₁₋₆alkyl, substituted C₁₋₆alkyl, C₂₋₆alkenyl, substituted C₂₋₆alkenyl, C₂₋₆alkynyl, substituted C₂₋₆alkynyl, C₃₋₆cycloalkyl, substituted C₃₋₆cycloalkyl, aryl, substituted aryl, and C₅₋₆heteroaryl, and substituted C₅₋₆heteroaryl; and

25 Y is selected from $-\text{CH}=\text{}$ and $-\text{N}=\text{}$.

DESCRIPTION OF THE INVENTION

Accordingly, it is an objective of certain embodiments of the invention to provide a compound that is effective in managing pain without causing the side effects mentioned above.

5 It is another objective of certain embodiments of the invention to provide a compound that is a CB₂ receptor ligand such as a CB₂ agonist that may be useful in managing pain and/or treating other symptoms or diseases.

Definitions

10 Unless specified otherwise within this specification, the nomenclature used in this specification generally follows the examples and rules stated in *Nomenclature of Organic Chemistry, Sections A, B, C, D, E, F, and H*, Pergamon Press, Oxford, 1979, which is incorporated by references herein for its exemplary chemical structure names and rules on naming chemical structures.

"CB₁/CB₂ receptors" means CB₁ and/or CB₂ receptors.

15 The term "C_{m-n}" or "C_{m-n} group" used alone or as a prefix, refers to any group having m to n carbon atoms.

The term "hydrocarbon" used alone or as a suffix or prefix, refers to any structure comprising only carbon and hydrogen atoms up to 14 carbon atoms.

20 The term "hydrocarbon radical" or "hydrocarbyl" used alone or as a suffix or prefix, refers to any structure as a result of removing one or more hydrogens from a hydrocarbon.

The term "alkyl" used alone or as a suffix or prefix, refers to monovalent straight or branched chain hydrocarbon radicals comprising 1 to about 12 carbon atoms. Unless otherwise specified, "alkyl" general includes both saturated alkyl and
25 unsaturated alkyl.

The term "alkylene" used alone or as suffix or prefix, refers to divalent straight or branched chain hydrocarbon radicals comprising 1 to about 12 carbon atoms, which serves to link two structures together.

30 The term "alkenyl" used alone or as suffix or prefix, refers to a monovalent straight or branched chain hydrocarbon radical having at least one carbon-carbon double bond and comprising at least 2 up to about 12 carbon atoms.

The term "alkynyl" used alone or as suffix or prefix, refers to a monovalent straight or branched chain hydrocarbon radical having at least one carbon-carbon triple bond and comprising at least 2 up to about 12 carbon atoms.

5 The term "cycloalkyl," used alone or as suffix or prefix, refers to a monovalent ring-containing hydrocarbon radical comprising at least 3 up to about 12 carbon atoms.

The term "cycloalkenyl" used alone or as suffix or prefix, refers to a monovalent ring-containing hydrocarbon radical having at least one carbon-carbon double bond and comprising at least 3 up to about 12 carbon atoms.

10 The term "cycloalkynyl" used alone or as suffix or prefix, refers to a monovalent ring-containing hydrocarbon radical having at least one carbon-carbon triple bond and comprising about 7 up to about 12 carbon atoms.

The term "aryl" used alone or as suffix or prefix, refers to a monovalent hydrocarbon radical having one or more polyunsaturated carbon rings having
15 aromatic character, (e.g., $4n + 2$ delocalized electrons) and comprising 5 up to about 14 carbon atoms, wherein the radical is located on a carbon of the aromatic ring. For example, aryl may be selected from phenyl and naphthyl.

The term "non-aromatic group" or "non-aromatic" used alone, as suffix or as prefix, refers to a chemical group or radical that does not contain a ring having
20 aromatic character (e.g., $4n + 2$ delocalized electrons).

The term "arylene" used alone or as suffix or prefix, refers to a divalent hydrocarbon radical having one or more polyunsaturated carbon rings having aromatic character, (e.g., $4n + 2$ delocalized electrons) and comprising 5 up to about 14 carbon atoms, which serves to link two structures together.

25 The term "heterocycle" used alone or as a suffix or prefix, refers to a ring-containing structure or molecule having one or more multivalent heteroatoms, independently selected from N, O, P and S, as a part of the ring structure and including at least 3 and up to about 20 atoms in the ring(s). Heterocycle may be saturated or unsaturated, containing one or more double bonds, and heterocycle may
30 contain more than one ring. When a heterocycle contains more than one ring, the rings may be fused or unfused. Fused rings generally refer to at least two rings sharing two atoms therebetween. Heterocycle may have aromatic character or may not have aromatic character.

The term "heteroalkyl" used alone or as a suffix or prefix, refers to a radical formed as a result of replacing one or more carbon atoms of an alkyl with one or more heteroatoms selected from N, O, P and S.

5 The term "heteroaromatic" used alone or as a suffix or prefix, refers to a ring-containing structure or molecule having one or more multivalent heteroatoms, independently selected from N, O, P and S, as a part of the ring structure and including at least 3 and up to about 20 atoms in the ring(s), wherein the ring-containing structure or molecule has an aromatic character (*e.g.*, $4n + 2$ delocalized electrons).

0 The term "heterocyclic group," "heterocyclic moiety," "heterocyclic," or "heterocyclo" used alone or as a suffix or prefix, refers to a radical derived from a heterocycle by removing one or more hydrogens therefrom.

15 The term "heterocyclyl" used alone or as a suffix or prefix, refers a monovalent radical derived from a heterocycle by removing one hydrogen from a carbon of a ring of the heterocycle.

The term "heterocyclylene" used alone or as a suffix or prefix, refers to a divalent radical derived from a heterocycle by removing two hydrogens therefrom, which serves to link two structures together.

20 The term "heteroaryl" used alone or as a suffix or prefix, refers to a heterocyclyl having aromatic character, wherein the radical of the heterocyclyl is located on a carbon of an aromatic ring of the heterocyclyl.

The term "heterocylcoalkyl" used alone or as a suffix or prefix, refers to a heterocyclyl that does not have aromatic character.

25 The term "heteroarylene" used alone or as a suffix or prefix, refers to a heterocyclylene having aromatic character.

The term "heterocycloalkylene" used alone or as a suffix or prefix, refers to a heterocyclylene that does not have aromatic character.

The term "six-membered" used as prefix refers to a group having a ring that contains six ring atoms.

30 The term "five-membered" used as prefix refers to a group having a ring that contains five ring atoms.

A five-membered ring heteroaryl is a heteroaryl with a ring having five ring atoms wherein 1, 2 or 3 ring atoms are independently selected from N, O and S.

Exemplary five-membered ring heteroaryls are thienyl, furyl, pyrrolyl, imidazolyl, thiazolyl, oxazolyl, pyrazolyl, isothiazolyl, isoxazolyl, 1,2,3-triazolyl, tetrazolyl, 1,2,3-thiadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-triazolyl, 1,2,4-thiadiazolyl, 1,2,4-oxadiazolyl, 1,3,4-triazolyl, 1,3,4-thiadiazolyl, and 1,3,4-oxadiazolyl.

5 A six-membered ring heteroaryl is a heteroaryl with a ring having six ring atoms wherein 1, 2 or 3 ring atoms are independently selected from N, O and S.

Exemplary six-membered ring heteroaryls are pyridyl, pyrazinyl, pyrimidinyl, triazinyl and pyridazinyl.

10 The term "substituted" used as a prefix refers to a structure, molecule or group, wherein one or more hydrogens are replaced with one or more C₁₋₁₂hydrocarbon groups, or one or more chemical groups containing one or more heteroatoms selected from N, O, S, F, Cl, Br, I, and P. Exemplary chemical groups containing one or more heteroatoms include heterocyclyl, -NO₂, -OR, -Cl, -Br, -I, -F, -CF₃, -C(=O)R, -C(=O)OH, -NH₂, -SH, -NHR, -NR₂, -SR, -SO₃H, -SO₂R, -S(=O)R, -CN, -OH, -C(=O)OR, -C(=O)NR₂, -NRC(=O)R, oxo (=O), imino (=NR), thio (=S), and oximino (=N-OR), wherein each "R" is a C₁₋₁₂hydrocarbyl. For example, substituted phenyl may refer to nitrophenyl, pyridylphenyl, methoxyphenyl, chlorophenyl, aminophenyl, etc., wherein the nitro, pyridyl, methoxy, chloro, and amino groups may replace any suitable hydrogen on the phenyl ring.

20 Ring nitrogen atoms of five-membered heteroaryl, heterocyclyl or bicyclic heteroaryl may be unsubstituted or substituted, if such a substitution is chemically possible without quaternization of said ring nitrogen, preferably with moieties independently selected from the group consisting of C₁₋₆alkyl, and -C(=O)R, wherein R is a C₁₋₆alkyl.

25 The term "substituted" used as a suffix of a first structure, molecule or group, followed by one or more names of chemical groups refers to a second structure, molecule or group, which is a result of replacing one or more hydrogens of the first structure, molecule or group with the one or more named chemical groups. For example, a "phenyl substituted by nitro" refers to nitrophenyl.

30 The term "optionally substituted" refers to both groups, structures, or molecules that are substituted and those that are not substituted.

Heterocycle includes, for example, monocyclic heterocycles such as: aziridine, oxirane, thiirane, azetidine, oxetane, thietane, pyrrolidine, pyrroline,

imidazolidine, pyrazolidine, pyrazoline, dioxolane, sulfolane 2,3-dihydrofuran, 2,5-dihydrofuran tetrahydrofuran, thiophane, piperidine, 1,2,3,6-tetrahydro-pyridine, piperazine, morpholine, thiomorpholine, pyran, thiopyran, 2,3-dihydropyran, tetrahydropyran, 1,4-dihydropyridine, 1,4-dioxane, 1,3-dioxane, dioxane,
5 homopiperidine, 2,3,4,7-tetrahydro-1*H*-azepine homopiperazine, 1,3-dioxepane, 4,7-dihydro-1,3-dioxepin, and hexamethylene oxide.

In addition, heterocycle includes aromatic heterocycles, for example, pyridine, pyrazine, pyrimidine, pyridazine, thiophene, furan, furazan, pyrrole, imidazole, thiazole, oxazole, pyrazole, isothiazole, isoxazole, 1,2,3-triazole, tetrazole, 1,2,3-
10 thiadiazole, 1,2,3-oxadiazole, 1,2,4-triazole, 1,2,4-thiadiazole, 1,2,4-oxadiazole, 1,3,4-triazole, 1,3,4-thiadiazole, and 1,3,4- oxadiazole.

Additionally, heterocycle encompass polycyclic heterocycles, for example, indole, indoline, isoindoline, quinoline, tetrahydroquinoline, isoquinoline, tetrahydroisoquinoline, 1,4-benzodioxan, coumarin, dihydrocoumarin, benzofuran,
15 2,3-dihydrobenzofuran, isobenzofuran, chromene, chroman, isochroman, xanthene, phenoxathiin, thianthrene, indolizine, isoindole, indazole, purine, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, phenanthridine, perimidine, phenanthroline, phenazine, phenothiazine, phenoxazine, 1,2-benzisoxazole, benzothiophene, benzoxazole, benzthiazole, benzimidazole,
20 benzotriazole, thioxanthine, carbazole, carboline, acridine, pyrolizidine, and quinolizidine.

One type of polycyclic heterocycles is bicyclic heteroaromatic ring system. A bicyclic heteroaromatic ring system is a ring system having two five- or six-membered heteroaromatic rings, or a phenyl and a five- or six-membered
25 heteroaromatic ring, or a phenyl and a heterocyclyl ring, or a five- or six-membered heteroaromatic ring and a heterocyclyl ring, connected by a ring fusion, said bicyclic heteroaromatic ring system comprising 8 to 12 ring atoms wherein 1, 2 or 3 of the ring atoms are independently selected from N, O and S.

For example, bicyclic heteroaromatic ring systems may be selected from
30 indole, indoline, quinoline, tetrahydroquinoline, isoquinoline, tetrahydroisoquinoline, 1,4-benzodioxan, coumarin, dihydrocoumarin, benzofuran, 2,3-dihydrobenzofuran, 1,2-benzisoxazole, benzothiophene, benzoxazole, benzthiazole, benzimidazole, benzotriazole, pyrolizidine, and quinolizidine.

In addition to the polycyclic heterocycles described above, heterocycle includes polycyclic heterocycles wherein the ring fusion between two or more rings includes more than one bond common to both rings and more than two atoms common to both rings. Examples of such bridged heterocycles include quinuclidine, diazabicyclo[2.2.1]heptane and 7-oxabicyclo[2.2.1]heptane.

Heterocyclyl includes, for example, monocyclic heterocyclyls, such as: aziridinyl, oxiranyl, thiranyl, azetidiny, oxetanyl, thietanyl, pyrrolidinyl, pyrrolinyl, imidazolidinyl, pyrazolidinyl, pyrazolinyl, dioxolanyl, sulfolanyl, 2,3-dihydrofuranyl, 2,5-dihydrofuranyl, tetrahydrofuranyl, thiophanyl, piperidinyl, 1,2,3,6-tetrahydro-
pyridinyl, piperazinyl, morpholinyl, thiomorpholinyl, pyranyl, thiopyranyl, 2,3-dihydropyranyl, tetrahydropyranyl, 1,4-dihydropyridinyl, 1,4-dioxanyl, 1,3-dioxanyl, dioxanyl, homopiperidinyl, 2,3,4,7-tetrahydro-1*H*-azepinyl, homopiperazinyl, 1,3-dioxepanyl, 4,7-dihydro-1,3-dioxepinyl, and hexamethylene oxidyl.

In addition, heterocyclyl includes aromatic heterocyclyls or heteroaryl, for example, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, thienyl, furyl, furazanyl, pyrrolyl, imidazolyl, thiazolyl, oxazolyl, pyrazolyl, isothiazolyl, isoxazolyl, 1,2,3-triazolyl, tetrazolyl, 1,2,3-thiadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-triazolyl, 1,2,4-thiadiazolyl, 1,2,4-oxadiazolyl, 1,3,4-triazolyl, 1,3,4-thiadiazolyl, and 1,3,4 oxadiazolyl.

Additionally, heterocyclyl encompasses polycyclic heterocyclyls (including both aromatic or non-aromatic), for example, indolyl, indolinyl, isoindolinyl, quinolinyl, tetrahydroquinolinyl, isoquinolinyl, tetrahydroisoquinolinyl, 1,4-benzodioxanyl, coumarinyl, dihydrocoumarinyl, benzofuranyl, 2,3-dihydrobenzofuranyl, isobenzofuranyl, chromenyl, chromanyl, isochromanyl, xanthenyl, phenoxathiinyl, thianthrenyl, indoliziny, isoindolyl, indazolyl, purinyl, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnolinyl, pteridinyl, phenanthridinyl, perimidinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, phenoxazinyl, 1,2-benzisoxazolyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benzimidazolyl, benztriazolyl, thioxanthinyl, carbazolyl, carbolinyl, acridinyl, pyrolizidinyl, and quinolizidinyl.

In addition to the polycyclic heterocyclyls described above, heterocyclyl includes polycyclic heterocyclyls wherein the ring fusion between two or more rings includes more than one bond common to both rings and more than two atoms

common to both rings. Examples of such bridged heterocycles include quinuclidinyl, diazabicyclo[2.2.1]heptyl; and 7-oxabicyclo[2.2.1]heptyl.

The term "alkoxy" used alone or as a suffix or prefix, refers to radicals of the general formula $-O-R$, wherein $-R$ is selected from a hydrocarbon radical. Exemplary alkoxy includes methoxy, ethoxy, propoxy, isopropoxy, butoxy, t-butoxy, isobutoxy, cyclopropylmethoxy, allyloxy, and propargyloxy.

The term "aryloxy" used alone or as suffix or prefix, refers to radicals of the general formula $-O-Ar$, wherein $-Ar$ is an aryl.

The term "heteroaryloxy" used alone or as suffix or prefix, refers to radicals of the general formula $-O-Ar'$, wherein $-Ar'$ is a heteroaryl.

The term "amine" or "amino" used alone or as a suffix or prefix, refers to radicals of the general formula $-NRR'$, wherein R and R' are independently selected from hydrogen or a hydrocarbon radical.

"Acyl" used alone, as a prefix or suffix, means $-C(=O)-R$, wherein $-R$ is an optionally substituted hydrocarbyl, hydrogen, amino or alkoxy. Acyl groups include, for example, acetyl, propionyl, benzoyl, phenyl acetyl, carboethoxy, and dimethylcarbamoyl.

Halogen includes fluorine, chlorine, bromine and iodine.

"Halogenated," used as a prefix of a group, means one or more hydrogens on the group are replaced with one or more halogens.

The term "electron-withdrawing group" refers to a chemical group has an electronegativity greater than a methyl group. Electronegativity measures the tendency of a group or atom to attract electrons. Exemplary electron-withdrawing groups are $-CH_2CF_3$, $-CF_3$, $-F$, $-Cl$, $-Br$, $-NO_2$, $-CN$, $-OH$, $-CHO$, $-C(=O)-R'$, $-OR'$ and C_1 -hydrocarbyl substituted by one or more groups selected from $-F$, $-Cl$, $-Br$, $-NO_2$, $-CN$, $-OH$, $-CHO$, $-C(=O)-R'$ and $-OR'$, wherein R' is a C_{1-3} alkyl.

"RT" or "rt" means room temperature.

A first ring group being "fused" with a second ring group means the first ring and the second ring share at least two atoms therebetween.

"Link," "linked," or "linking," unless otherwise specified, means covalently linked or bonded.

When a first group, structure, or atom is "directly connected" to a second group, structure or atom, at least one atom of the first group, structure or atom forms a chemical bond with at least one atom of the second group, structure or atom.


"Saturated carbon" means a carbon atom in a structure, molecule or group wherein all the bonds connected to this carbon atom are single bond. In other words, there is no double or triple bonds connected to this carbon atom and this carbon atom generally adopts an sp^3 atomic orbital hybridization.

"Unsaturated carbon" means a carbon atom in a structure, molecule or group wherein at least one bond connected to this carbon atom is not a single bond. In other words, there is at least one double or triple bond connected to this carbon atom and this carbon atom generally adopts an sp or sp^2 atomic orbital hybridization.

The term "therapy" also includes "prophylaxis" unless there are specific indications to the contrary. The term "therapeutic" and "therapeutically" should be construed accordingly. The term "therapy" within the context of the present invention further encompasses to administer an effective amount of a compound of the present invention, to mitigate either a pre-existing disease state, acute or chronic, or a recurring condition. This definition also encompasses prophylactic therapies for prevention of recurring conditions and continued therapy for chronic disorders.

20 Description of Embodiments

In one embodiment, the invention provides compounds represented by the formula I, pharmaceutically acceptable salts thereof, diastereomers, enantiomers and mixtures thereof:



25 wherein

R^{F1} and R^{F2} are independently C_{1-10} electron-withdrawing groups;

Z is selected from O= and S=;

R^1 is selected from C_{1-10} alkyl; C_{1-10} alkyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; C_{2-10} alkenyl; C_{2-10} alkenyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; C_{2-10} alkynyl; C_{2-10} alkynyl

substituted by at least one of halogen, cyano, acetoxymethyl and nitro; R^3R^4N - C_{1-6} alkyl; $R^3R^4NC(=O)-C_{1-6}$ alkyl; R^3O-C_{1-6} alkyl; $R^3OC(=O)-C_{1-6}$ alkyl; $R^3C(=O)-C_{1-6}$ alkyl; $R^3C(=O)NR^3-C_{1-6}$ alkyl; $R^3R^4NSO_2-C_{1-6}$ alkyl; $R^3CSO_2N(R^4)-C_{1-6}$ alkyl; $R^3R^4NC(=O)N(R^5)-C_{1-6}$ alkyl; $R^3R^4NSO_2N(R^5)-C_{1-6}$ alkyl; aryl- C_{1-6} alkyl; aryl- $C(=O)-C_{1-6}$ alkyl; heterocyclyl- C_{1-6} alkyl; heterocyclyl- $C(=O)-C_{1-6}$ alkyl; substituted aryl- C_{1-6} alkyl; substituted aryl- $C(=O)-C_{1-6}$ alkyl; substituted heterocyclyl- C_{1-6} alkyl; substituted heterocyclyl- $C(=O)-C_{1-6}$ alkyl; and C_{1-10} hydrocarbylamino;

R^3 , R^4 and R^5 are independently selected from -H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and a divalent C_{1-6} group that together with another divalent C_{1-6} group forms a portion of a ring;

X is a C_{1-10} divalent group that separates groups connected thereto by one or two atoms;

Ar is a C_{4-12} divalent aromatic group;

R^2 is selected from C_{1-6} alkyl, substituted C_{1-6} alkyl, C_{2-6} alkenyl, substituted C_{2-6} alkenyl, C_{2-6} alkynyl, substituted C_{2-6} alkynyl, C_{3-6} cycloalkyl, substituted C_{3-6} cycloalkyl, aryl, substituted aryl, and C_{5-6} heteroaryl, and substituted C_{5-6} heteroaryl; and

Y is selected from $-CH=$ and $-N=$.

In another embodiment, compounds of the present invention may be those of formula I, wherein

R^{F1} and R^{F2} are independently C_{1-6} alkyl substituted by one or more groups selected from -F, -Cl, -Br, $-NO_2$, -CN, -OH, -CHO, $-C(=O)-R'$ and $-OR'$, wherein R' is a C_{1-3} alkyl;

R^1 is selected from C_{1-8} alkyl; C_{2-8} alkenyl; C_{2-8} alkynyl; aryl- C_{1-6} alkyl; aryl- C_{1-6} alkyl with the aryl substituted by at least one group selected from C_{1-6} alkyl, acetoxymethyl, nitro and halogen; $R^8R^9NC_{1-6}$ alkyl; R^8OC_{1-6} alkyl; cycloalkyl- C_{1-6} alkyl; heterocycloalkyl- C_{1-6} alkyl; heterocycloalkyl- C_{1-6} alkyl with the heterocycloalkyl thereof substituted by at least one group selected from C_{1-8} alkyl, acetoxymethyl, nitro and halogen; C_{1-6} alkylaryl; C_{1-6} alkyl- $C(=O)-$; C_{6-8} aryl- $C(=O)-$; C_{4-8} heteroaryl- $C(=O)-$; heteroaryl- C_{1-6} alkyl; heteroaryl- C_{1-6} alkyl with the heteroaryl thereof substituted by at least one group selected from C_{1-6} alkyl, acetoxymethyl, nitro and halogen; and R^NC_{1-6} alkyl;

R^N is an oxidized pyridyl wherein the nitrogen atom of the pyridyl ring is in an oxidized state (N^+-O^-);

R^2 is selected from C_{1-6} alkyl, C_{1-6} alkyl substituted by at least one fluorine, C_{2-6} alkenyl, C_{2-6} alkenyl substituted by at least one fluorine, C_{2-6} alkynyl, C_{2-6} alkynyl substituted by at least one fluorine, C_{3-6} cycloalkyl, substituted C_{3-6} cycloalkyl, aryl, substituted aryl, and C_{5-6} heteroaryl, and substituted C_{5-6} heteroaryl;

R^3 , R^4 and R^5 are independently selected from -H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and a divalent C_{1-6} group that together with another divalent C_{1-6} group forms a portion of a ring; and

X is selected from $-NR^6-$, $-C(=O)-$, $-CH_2-CH_2-$, $-CH=CH-$, $-O-$, $-C(R^6)(R^7)-$, and $-S(O)_q-$, wherein q is 0, 1 or 2, wherein R^6 and R^7 are independently C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, OH, or H.

In a further embodiment, compounds of the present invention may be those of formula I, wherein

R^{F1} and R^{F2} are independently selected from $-CF_3$, $-CH_2CF_3$, $-CH_2CHF_2$, $-CHFCH_2F$, $-CHFCHF_2$, $-CHFCH_2F$, $-CF_2CF_3$, $-CF_2CH_3$, $-CF_2CH_2F$, $-CF_2CHF_2$, $-CCl_3$, $-CH_2CCl_3$, $-CH_2CHCl_2$, $-CH_2CBr_3$, $-CH_2CHBr_2$, $-CH_2NO_2$, $-CH_2CH_2NO_2$, $-CH_2CN$, $-CH_2CH_2CN$, and $-CH_2CH_2OCH_3$;

R^1 is selected from C_{1-8} alkyl; C_{2-8} alkenyl; C_{2-8} alkynyl; aryl- C_{1-6} alkyl; aryl- C_{1-6} alkyl with the aryl substituted by at least one group selected from C_{1-6} alkyl, acetoxymethyl, nitro and halogen; $R^8R^9NC_{1-6}$ alkyl; R^8OC_{1-6} alkyl; cycloalkyl- C_{1-6} alkyl; heterocycloalkyl- C_{1-6} alkyl; heterocycloalkyl- C_{1-6} alkyl with the heterocycloalkyl thereof substituted by at least one group selected from C_{1-8} alkyl, acetoxymethyl, nitro and halogen; C_{1-6} alkylaryl; C_{1-6} alkyl- $C(=O)-$; C_{6-8} aryl- $C(=O)-$; C_{4-8} heteroaryl- $C(=O)-$; heteroaryl- C_{1-6} alkyl; heteroaryl- C_{1-6} alkyl with the heteroaryl thereof substituted by at least one group selected from C_{1-6} alkyl, acetoxymethyl, nitro and halogen; and $R^N C_{1-6}$ alkyl;

R^N is an oxidized pyridyl wherein the nitrogen atom of the pyridyl ring is in an oxidized state (N^+-O^-);

Z is $O=$;

R^2 is selected from $-CH_3$, $-CH_2CH_3$, $-CH(CH_3)_2$, C_{3-6} cycloalkyl, $-CH_2CF_3$, $-CHF_2$, $-CF_3$ and aryl;

Ar is selected from an arylene; an heteroarylene; an arylene substituted by at least one group selected from C₁₋₆alkyl, halogen, trifluoromethyl, cyano, nitro, hydroxy and C₁₋₆alkoxy; and an heteroarylene substituted by at least one group selected from C₁₋₆alkyl, halogen, trifluoromethyl, cyano, nitro, hydroxy and C₁₋₆alkoxy, even more particularly, the arylene is *para*-arylene; and the heteroarylene is selected from the group consisting of six-membered ring *para*-heteroarylene and five-membered ring *meta*-heteroarylene; and

R⁸ and R⁹ are independently selected from the group consisting of -H and C₁₋₆alkyl.

In an even further embodiment, compounds of the present invention may be those of formula I, wherein

R^{F1} and R^{F2} are independently C₁₋₆ groups that comprise at least 30% fluorine by weight;

Z is O=;

R¹ is selected from ethyl, propyl, allyl, isopentyl, benzyl, dimethylaminoethyl, 4-pyridylmethyl, 2-pyridylmethyl, 1-pyrrolylethyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, 2-pyrrolidylmethyl, 3-pyrrolidylmethyl, N-methyl-2-pyrrolidylmethyl, N-methyl-3-pyrrolidylmethyl, 2-piperidylmethyl, 3-piperidylmethyl, 4-piperidylmethyl, N-methyl-2-piperidylmethyl, N-methyl-3-piperidylmethyl, N-methyl-4-piperidylmethyl, 3-thienylmethyl, 2-tetrahydrofuranylmethyl, 3-tetrahydrofuranylmethyl, 2-tetrahydropyranylmethyl, 3-tetrahydropyranylmethyl, 4-tetrahydropyranylmethyl, (2-nitrothiophene-5-yl)methyl, (1-methyl-1H-imidazole-2-yl)methyl, (5-(acetoxymethyl)-2-furanyl)methyl, (2,3-dihydro-1H-isoindeole-1-yl)methyl, and 5-(2-methylthiazolyl);

R² is selected from -CH₃, -CH₂CH₃, -CH(CH₃)₂, -CH₂CF₃, CF₃, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and phenyl;

X is selected from -CH₂- and -CH(CH₃)-; and

Ar is selected from a *para*-arylene; a *para*-arylene substituted with C₁₋₆alkyl, halogen, trifluoromethyl, cyano, nitro, hydroxy and C₁₋₆alkoxy; a six-membered ring *para*-heteroarylene; and a six-membered ring *para*-heteroarylene substituted with a group selected from C₁₋₆alkyl, halogen, trifluoromethyl, cyano, nitro, hydroxy and C₁₋₆alkoxy.

In an even further embodiment, R^{F1} and R^{F2} are $-\text{CH}_2\text{CF}_3$; R^2 is $-\text{CH}_2\text{CH}_3$; and Ar is selected from the group consisting of *para*-phenylene and *para*-pyridylene.

It will be understood that when compounds of the present invention contain one or more chiral centers, the compounds of the invention may exist in, and be
5 isolated as, enantiomeric or diastereomeric forms, or as a racemic mixture. The present invention includes any possible enantiomers, diastereomers, racemates or mixtures thereof, of a compound of Formula I. The optically active forms of the compound of the invention may be prepared, for example, by chiral chromatographic separation of a racemate, by synthesis from optically active starting materials or by
10 asymmetric synthesis based on the procedures described thereafter.

It will also be appreciated that certain compounds of the present invention may exist as geometrical isomers, for example E and Z isomers of alkenes. The present invention includes any geometrical isomer of a compound of Formula I. It will further be understood that the present invention encompasses tautomers of the
15 compounds of the formula I.

It will also be understood that certain compounds of the present invention may exist in solvated, for example hydrated, as well as unsolvated forms. It will further be understood that the present invention encompasses all such solvated forms of the compounds of the formula I.

20 Within the scope of the invention are also salts of the compounds of the formula I. Generally, pharmaceutically acceptable salts of compounds of the present invention may be obtained using standard procedures well known in the art, for example by reacting a sufficiently basic compound, for example an alkyl amine with a suitable acid, for example, HCl or acetic acid, to afford a physiologically acceptable
25 anion. It may also be possible to make a corresponding alkali metal (such as sodium, potassium, or lithium) or an alkaline earth metal (such as a calcium) salt by treating a compound of the present invention having a suitably acidic proton, such as a carboxylic acid or a phenol with one equivalent of an alkali metal or alkaline earth metal hydroxide or alkoxide (such as the ethoxide or methoxide), or a suitably basic
30 organic amine (such as choline or meglumine) in an aqueous medium, followed by conventional purification techniques.

In one embodiment, the compound of formula I above may be converted to a pharmaceutically acceptable salt or solvate thereof, particularly, an acid addition salt

such as a hydrochloride, hydrobromide, phosphate, acetate, fumarate, maleate, tartrate, citrate, methanesulphonate or *p*-toluenesulphonate.

The compounds of the invention may have activity as pharmaceuticals, in particular as modulators or ligands such as agonists, partial agonists, inverse agonist
5 or antagonists of CB₂ receptors. In one embodiment, the compounds of the invention may exhibit selective activity as agonists at CB₂ receptors, and are useful in the relief of pain, particularly chronic pain, e.g., chronic inflammatory pain, neuropathic pain, back pain, cancer pain and visceral pain. Compounds of the present invention may also be useful in treating acute pain. Additionally, compounds of the present
10 invention may be useful in other disease states in which degeneration or dysfunction of CB₂ receptors is present or implicated.

Thus, the invention provides a compound of formula I, or pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined for use in therapy.

In a further aspect, the present invention provides the use of a compound of
15 formula I, or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in the manufacture of a medicament for use in therapy.

The compounds of the present invention may be useful in therapy, especially for the therapy of various pain conditions including, but not limited to: acute pain, chronic pain, neuropathic pain, back pain, cancer pain, and visceral pain.

20 The compounds of the present invention may also be used in treating pain, cancer, multiple sclerosis, Parkinson's disease, transplant rejection, Huntington's chorea and/or Alzheimer's disease.

In use for therapy in a warm-blooded animal such as a human, the compound of the invention may be administered in the form of a conventional pharmaceutical
25 composition by any route including orally, intramuscularly, subcutaneously, topically, intranasally, intraperitoneally, intrathoracically, intravenously, epidurally, intrathecally, intracerebroventricularly and by injection into the joints.

In one embodiment of the invention, the route of administration may be oral, intravenous or intramuscular.

30 The dosage will depend on the route of administration, the severity of the disease, age and weight of the patient and other factors normally considered by the attending physician, when determining the individual regimen and dosage level at the most appropriate for a particular patient.

For preparing pharmaceutical compositions from the compounds of this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets, and suppositories.

5 A solid carrier can be one or more substances, which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, or table disintegrating agents; it can also be an encapsulating material.

10 In powders, the carrier is a finely divided solid, which is in a mixture with the finely divided compound of the invention, or the active component. In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

15 For preparing suppository compositions, a low-melting wax such as a mixture of fatty acid glycerides and cocoa butter is first melted and the active ingredient is dispersed therein by, for example, stirring. The molten homogeneous mixture is then poured into convenient sized moulds and allowed to cool and solidify.

Suitable carriers are magnesium carbonate, magnesium stearate, talc, lactose, sugar, pectin, dextrin, starch, tragacanth, methyl cellulose, sodium carboxymethyl cellulose, a low-melting wax, cocoa butter, and the like.

20 The term composition is also intended to include the formulation of the active component with encapsulating material as a carrier providing a capsule in which the active component (with or without other carriers) is surrounded by a carrier which is thus in association with it. Similarly, cachets are included.

Tablets, powders, cachets, and capsules can be used as solid dosage forms suitable for oral administration.

25 Liquid form compositions include solutions, suspensions, and emulsions. For example, sterile water or water propylene glycol solutions of the active compounds may be liquid preparations suitable for parenteral administration. Liquid compositions can also be formulated in solution in aqueous polyethylene glycol solution.

30 Aqueous solutions for oral administration can be prepared by dissolving the active component in water and adding suitable colorants, flavoring agents, stabilizers, and thickening agents as desired. Aqueous suspensions for oral use can be made by dispersing the finely divided active component in water together with a viscous

material such as natural synthetic gums, resins, methyl cellulose, sodium carboxymethyl cellulose, and other suspending agents known to the pharmaceutical formulation art.

Depending on the mode of administration, the pharmaceutical composition
5 will preferably include from 0.05% to 99%w (per cent by weight), more preferably from 0.10 to 50%w, of the compound of the invention, all percentages by weight being based on total composition.

A therapeutically effective amount for the practice of the present invention may be determined, by the use of known criteria including the age, weight and
10 response of the individual patient, and interpreted within the context of the disease which is being treated or which is being prevented, by one of ordinary skill in the art.

Within the scope of the invention is the use of any compound of formula I as defined above for the manufacture of a medicament.

Also within the scope of the invention is the use of any compound of formula I
15 for the manufacture of a medicament for the therapy of pain.

Additionally provided is the use of any compound according to Formula I for the manufacture of a medicament for the therapy of various pain conditions including, but not limited to: acute pain, chronic pain, neuropathic pain, acute pain, back pain, cancer pain, and visceral pain.

20 A further aspect of the invention is a method for therapy of a subject suffering from any of the conditions discussed above, whereby an effective amount of a compound according to the formula I above, is administered to a patient in need of such therapy.

Additionally, there is provided a pharmaceutical composition comprising a
25 compound of Formula I, or a pharmaceutically acceptable salt thereof, in association with a pharmaceutically acceptable carrier.

Particularly, there is provided a pharmaceutical composition comprising a compound of Formula I, or a pharmaceutically acceptable salt thereof, in association with a pharmaceutically acceptable carrier for therapy, more particularly for therapy
30 of pain.

Further, there is provided a pharmaceutical composition comprising a compound of Formula I, or a pharmaceutically acceptable salt thereof, in association

with a pharmaceutically acceptable carrier use in any of the conditions discussed above.

In one embodiment, the compounds of the invention are found to be active towards CB₂ receptors in warm-blooded animals, e.g., human. Particularly the compounds of the invention are found to be effective CB₂ receptor agonists. *In vitro* assays, *infra*, demonstrate these activities. In these *in vitro* assays, a compound is tested for their activity toward CB₂ receptors and dissociation constant (K_i) is obtained to determine the selective activity for a particular compound towards CB₂ receptors by measuring IC₅₀ of the compound. In the current context, IC₅₀ generally refers to the concentration of the compound at which 50% displacement of a standard radioactive CB₂ receptor ligand has been observed. Generally, a lower K_i for a particular compound towards CB₂ receptors means that the particular compound is a stronger ligand towards the CB₂ receptors. As a result, compounds with relatively low K_i towards CB₂ receptors are relatively strong CB₂ receptor ligands or potent CB₂ receptor agonists.

The activities of the compound towards CB₁ receptors are also measured in a similar assay. It is known that even though CB₁ receptor agonists are also effective in relieving or managing pains, their use is often associated with undesired CNS side-effects, e.g., psychoactive side effects and abuse potential. In one embodiment, we surprisingly find the particular compounds of the present invention tend to bind weakly to CB₁ receptors, have relatively high K_i's towards particular CB₁ receptors and thus low binding affinity toward CB₁ receptors. As a result, the compounds may show no or less side effects in treating pain in comparison with conventional cannabinoids.

CB₂ receptors are generally expressed in peripheral tissues, particularly in the immuno-competent and inflammatory cells, which are believed to have therapeutic effects related to immunomodulation, inflammation and bronchial constriction. The CB₂ receptor selective ligands such as CB₂ receptor agonists may have therapeutic utility in controlling diseases associated with these conditions.

It is substantially established that CB₂ receptor ligands (e.g. agonists) are generally effective in treating or managing pain in warm-blooded animals with little side effects such as addiction and the CB₂ receptor ligands may have similar effects on humans. For example, CB₂ receptors have been found to be expressed in

peritoneal mast cells which play a crucial role in amplifying the key nerve growth factor (NGF) which may induce inflammatory hyperalgesia. CB₂ receptors have also been shown to be involved in the attenuation of nitric oxide production from macrophages upon treatment of LPS.

5 In addition, *in vivo* pain models have shown that CB₂ receptor agonists may induce analgesia. In particular, it has been found that the expression of the CB₂ receptors is induced under conditions of immune cell activation and that a CB₂ agonist elicits anti-inflammatory and peripheral analgesic activity. Moreover, it has been shown that CB₂ activation inhibits mechanical hyperalgesia associated with
10 nerve injury. Further, it has been demonstrated that peripheral CB₂ receptors may mediate antinociception in the rat. Consequentially, in one embodiment, the compounds that show a lower K_i in the *in vitro* assays disclosed in the invention may be suitable to be used in therapies such as pain relief or pain management.

 CB₂ receptor selective ligands such as CB₂ receptor agonists (sometimes also
15 called CB₂ selective cannabinoids) have been shown to induce apoptosis in glioma cells, in both *in vitro* and *in vivo* settings. Further evidence has supported the antiproliferative effects of CB₂ receptor agonists on breast and prostate cancer cells. CB₂ receptor agonists can also induce tumor suppression in a rat model in which rat glioma C6 cells are inoculated intracerebrally in the rats. Both clinical and non-
20 clinical studies also show that CB₂ receptor ligands may also have a positive effect on multiple sclerosis.


 CB₂ receptor agonists may also be used in other therapies related directly or indirectly to immunomodulation such as treating asthma, bronchitis, chronic obstructive pulmonary disease (COPD), reversible airway obstruction, adult
25 respiratory distress syndrome, psoriasis, rheumatoid arthritis, and allergy. CB₂ receptor agonists may also be effective in treating Parkinson's disease, Huntington's chorea and Alzheimer's disease, and Crohn's disease. CB₂ receptor agonists may also be useful in transplant rejection therapy.

 Therefore, in another aspect, the compounds of the invention may be useful
30 and effective in treating LPS, pain, multiple sclerosis, asthma, bronchitis, chronic obstructive pulmonary disease (COPD), reversible airway obstruction, adult respiratory distress syndrome, psoriasis, rheumatoid arthritis, allergy, Parkinson's

disease, Huntington's chorea and Alzheimer's disease, Crohn's disease, and transplant rejection.

In a further aspect, the present invention provides a method of preparing a compound of formula I.

- 5 In one embodiment, the method of preparing the compound of the invention includes the step of reacting a compound represented by formula II with $R^2OArXC(=O)A$:



wherein

- 10 R^{F1} and R^{F2} are independently electron-withdrawing groups;
 Z is selected from O= and S=;
 R^1 is a C_{1-11} group;
 X is a divalent group that separates groups connected thereto by one or two atoms;
 15 A is selected from -OH, -Cl, -Br, and -I;
 Ar is selected from a divalent aromatic group;
 R^2 is a C_{1-6} group; and
 Y is selected from -CH= and -N=.
- In another embodiment, R^{F1} , R^{F2} , R^1 , R^2 , Z, X, Ar, and Y are the same as those
 20 defined in the context of defining formula I.


In a further embodiment, the reaction conditions for the above reaction step may be as follows:

- Process a) A compound of formula II and $R^2OArXC(=O)A$ may be reacted together in a suitable solvent (e.g. 1,2-dichloroethane) in the presence of a
 25 reducing agent (e.g., zinc) followed by treatment with an acid (e.g., HCl) with heating (e.g., at 70-100°C) to form a compound of formula I, wherein A is -Cl, -Br or -I.

Process b) A compound of formula II and $R^2OArXC(=O)OH$ may be reacted together in a suitable polar solvent (e.g., dimethylformamide (DMF)) in the

presence of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyl uronium hexafluorophosphate (HATU) and a base such as diisopropylethylamine (DIPEA) followed by treatment with an acid (e.g. HCl) with heating (e.g., at 70-100°C) to form a compound of formula I.

- 5 In another embodiment, the method of preparing the compound of the invention includes the step of reacting a compound represented by formula III with formaldehyde:



wherein


- 10 r and s are selected from 0, 1 and 2;
 R^{10} is selected from C_{1-6} alkylene, -O-, and $-NR^{11}-$, wherein R^{11} is a C_{1-6} alkyl;
 R^{F1} and R^{F2} are independently electron-withdrawing groups;
 X is a divalent group that separates groups connected thereto by one or two atoms;
 15 Ar is a divalent aromatic group;
 A is selected from -OH, -Cl, -Br, and -I.
 R^2 is a C_{1-6} group; and
 Y is selected from -CH= and -N=.
- In a further embodiment, R^{F1} , R^{F2} , R^2 , X, Ar, and Y are the same as those
 20 defined in the context of defining formula I.

In an even further embodiment, the reaction conditions for the above reacting step may be as follows:

- A compound of formula III and formaldehyde may be reacted together in a suitable solvent (e.g. methanol, or tetrahydrofuran (THF)) in the presence of a
 25 reducing or hydrogenation agent (e.g., $NaCNBH_3$, $NaBH(OAc)_3$), optionally further in the presence of an acid such as acetic acid, to form a compound of formula I.

In an even further embodiment, the compounds of the present invention can be prepared according to the synthetic routes as depicted in Schemes 1 and 2.


Scheme 1



$\text{R}^1, \text{R}^2, \text{Ar}, \text{X}$ are the same as defined above.

5

Scheme 2



Ar, X and R^2 are the same as defined above.

10 EXAMPLES

The invention will further be described in more detail by the following Examples which describe methods whereby compounds of the present invention may be prepared, purified, analyzed and biologically tested, and which are not to be construed as limiting the invention.

15

Example 1: Biological Evaluation

hCB₁ and hCB₂ receptor binding

Human CB₁ receptor from Receptor Biology (hCB₁) or human CB₂ receptor from BioSignal (hCB₂) membranes are thawed at 37 °C, passed 3 times through a 25-gauge blunt-end needle, diluted in the cannabinoid binding buffer (50 mM Tris, 2.5 mM EDTA, 5 mM MgCl₂, and 0.5 mg/mL BSA fatty acid free, pH 7.4) and aliquots containing the appropriate amount of protein are distributed in 96-well plates. The IC₅₀ of the compounds of the invention at hCB₁ and hCB₂ are evaluated from 10-point dose-response curves done with ³H-CP55,940 at 20000 to 25000 dpm per well (0.17-0.21 nM) in a final volume of 300 µl. The total and non-specific binding are determined in the absence and presence of 0.2 µM of HU210 respectively. The plates are vortexed and incubated for 60 minutes at room temperature, filtered through Unifilters GF/B (presoaked in 0.1% polyethyleneimine) with the Tomtec or Packard harvester using 3 mL of wash buffer (50 mM Tris, 5 mM MgCl₂, 0.5 mg BSA pH 7.0). The filters are dried for 1 hour at 55 °C. The radioactivity (cpm) is counted in a TopCount (Packard) after adding 65 µl/well of MS-20 scintillation liquid.

hCB₁ and hCB₂ GTPγS binding

Human CB₁ receptor from Receptor Biology (hCB₁) or human CB₂ receptor membranes (BioSignal) are thawed at 37 °C, passed 3 times through a 25-gauge blunt-end needle and diluted in the GTPγS binding buffer (50 mM Hepes, 20 mM NaOH, 100 mM NaCl, 1 mM EDTA, 5 mM MgCl₂, pH 7.4, 0.1% BSA). The EC₅₀ and E_{max} of the compounds of the invention are evaluated from 10-point dose-response curves done in 300µl with the appropriate amount of membrane protein and 100000-130000 dpm of GTPγ³⁵S per well (0.11 –0.14 nM). The basal and maximal stimulated binding is determined in absence and presence of 1 µM (hCB₂) or 10 µM (hCB₁) Win 55,212-2 respectively. The membranes are pre-incubated for 5 minutes with 56.25 µM (hCB₂) or 112.5 µM (hCB₁) GDP prior to distribution in plates (15 µM (hCB₂) or 30 µM (hCB₁) GDP final). The plates are vortexed and incubated for 60 minutes at room temperature, filtered on Unifilters GF/B (presoaked in water) with the Tomtec or Packard harvester using 3 ml of wash buffer (50 mM Tris, 5 mM MgCl₂, 50 mM NaCl, pH 7.0). The filters are dried for 1 hour at 55 °C. The radioactivity (cpm) is counted in a TopCount (Packard) after adding 65 µl/well of MS-20 scintillation liquid. Antagonist reversal studies are done in the same way

except that (a) an agonist dose-response curve is done in the presence of a constant concentration of antagonist, or (b) an antagonist dose-response curve is done in the presence of a constant concentration of agonist.

Based on the above assays, the dissociation constant (K_i) for a particular compound of the invention towards a particular receptor is determined using the following equation:

$$K_i = IC_{50} / (1 + [rad] / K_d),$$

Wherein IC_{50} is the concentration of the compound of the invention at which 50% displacement has been observed;

[rad] is a standard or reference radioactive ligand concentration at that moment; and



K_d is the dissociation constant of the radioactive ligand towards the particular receptor.







Biological data for certain compounds of the invention are listed in Table 1







below.







Table 1







Biological data







Compound No.		hCB1 Ki (nM)	hCB2 Ki (nM)
1		2799.9	3.1
2		5084.4	5.6





3		84.8	0.8
4		4694.1	13.9
5		5388.1	44.1
6		5531.5	42.7
7		5531.5	53.1
8		5350.0	16.7


9		5568.0	164.9
10		3502.4	28.2
11		3234.4	12.0
12		5715.4	12.5
13		565.4	5.4
14		5749.8	14.8

15	 Chiral	5749.8	9.6
16	 Chiral	1414.0	3.3
17	 Chiral	2964.0	19.3
18	 Chiral	5583.4	179.8
19	 Chiral	1315.5	7.3
20	 Chiral	2729.2	11.2

21		5433.4	15.4
22		5523.1	23.1
23		5572.3	55.8
24		5746.9	138.4
25		1555.4	2.0
26		5528.7	6.6

27		1356.8	3.2
28		5538.6	40.5
29		5814.9	68.7
30		454.9	3.1
31		826.4	13.5
32		142.0	2.7

33		5178.8	446.4
34		5378.2	78.2
35		5572.3	1585.1
36		5751.1	14.0


Example 2: synthesis of intermediates 1-39**5 Intermediate 1****4-Fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide**

4-Fluoro-3-nitrobenzoic acid (2.50 g, 13.5 mmol) was refluxed in a 2:1 mixture of $\text{CH}_2\text{Cl}_2/\text{SOCl}_2$ (150 mL) for 5h. The solvent was concentrated and the residue was dissolved in CH_2Cl_2 (50 mL). Another CH_2Cl_2 solution (50 mL) of diisopropylethylamine (DIPEA) (3.50 mL, 20.3 mmol) and bis(2,2,2-

trifluoroethyl)amine (4.90 g, 27.0 mmol) was then added dropwise to the cold stirring solution (0 °C) of the acid chloride. The solution was stirred at RT overnight. The solution was then washed with 5% KHSO₄ solution, saturated NaHCO₃ solution, and brine and dried over anhydrous MgSO₄. The product was purified by flash chromatography on silica gel using 3:1/HEX:EtOAc. Yield: 3.08 g (66%); ¹H NMR (CDCl₃) 4.17 (brs, 4H), 7.40 (t, J = 8.59Hz, 1H), 7.66 (m, 2H), 8.10 (d, J = 6.84Hz, 1H).

Intermediate 2


4-[(3-Methylbutyl)amino]-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)-benzamide



4-Fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (170 mg, 0.488 mmol) and isoamylamine (0.068 mL, 0.586 mmol) were stirred in 3 mL of EtOH containing Et₃N (0.136 mL, 0.976 mmol) at 80 °C for 3h. The solvent was concentrated. The residue was dissolved in EtOAc and washed with saturated NaHCO₃, brine and dried over anhydrous Na₂SO₄. The product was purified by flash chromatography on silica gel using 3:1 / hexanes:EtOAc as eluent affording intermediate 2 as a yellow oil. Yield: 203 mg (99%); ¹H NMR (CDCl₃) 1.00 (d, J = 6.64Hz, 6H), 1.67 (m, 2H), 1.79 (m, 1H), 3.36 (m, 2H), 4.25 (q, J = 8.59Hz, 4H), 6.95 (d, J = 8.98Hz, 1H), 7.57 (d, J = 8.98Hz, 1H), 8.30 (m, 2H); MS (ESI) 416.22 (MH⁺).

Intermediate 3


4-[(Cyclopropylmethyl)amino]-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)-benzamide



Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (100 mg, 0.287 mmol) and cyclopropylmethyl amine (0.030 mL, 0.344 mmol) in 3 mL of EtOH containing Et₃N (0.080 mL, 0.574 mmol). The product was purified by flash chromatography on silica gel using 4:1 /
 5 hexanes:EtOAc as eluent. Yield: 91 mg (80%); ¹H NMR (CDCl₃) 0.35 (m, 2H), 0.70 (m, 2H), 1.21 (m, 1H), 3.21 (m, 2H), 4.25 (q, *J* = 8.59Hz, 4H), 6.91 (d, *J* = 8.98Hz, 1H), 7.54 (d, *J* = 8.79Hz, 1H), 8.30 (s, 1H), 8.35 (s, 1H); MS (ESI) 400.19 (MH⁺).

Intermediate 4

10 4-[(Cyclohexylmethyl)amino]-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)-benzamide




Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (100 mg, 0.287 mmol) and cyclohexylmethyl amine (0.041 mL, 0.315 mmol) in 3 mL of EtOH containing Et₃N (0.080 mL, 0.574
 15 mmol). The product was purified by flash chromatography on silica gel using 4:1 / hexanes:EtOAc as eluent. Yield: 107 mg (85%); ¹H NMR (CDCl₃) 1.05 (m, 2H), 1.24 (m, 4H), 1.69 (m, 2H), 1.75 (m, 2H), 1.83 (m, 1H), 3.17 (t, *J* = 6.64Hz, 2H), 4.23 (q, *J* = 8.53Hz, 4H), 6.91 (d, *J* = 8.98Hz, 1H), 7.51 (d, *J* = 8.89Hz, 1H), 8.28 (s, 1H), 8.39 (s, 1H); MS (ESI) 442.29 (MH⁺).

20

Intermediate 5

4-[(2-Furanylmethyl)amino]-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)-benzamide




Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (100 mg, 0.287 mmol) and furfurylamine (0.028
 25

mL, 0.315 mmol) in 3 mL of EtOH containing Et₃N (0.080 mL, 0.574 mmol). The product was purified by flash chromatography on silica gel using 4:1 / hexanes:EtOAc as eluent. Yield: 115mg (94%); ¹H NMR (CDCl₃) 4.23 (q, J = 8.40Hz, 4H), 4.56 (d, J = 5.66Hz, 2H), 6.31 (d, J = 3.12Hz, 1H), 6.35 (m, 1H), 7.04 (d, J = 8.98Hz, 1H), 7.39 (s, 1H), 7.53 (d, J = 8.89Hz, 1H), 8.28 (s, 1H), 8.55 (s, 1H); MS (ESI) 426.21 (MH⁺).

Intermediate 6

2-[[[4-[[Bis(2,2,2-trifluoroethyl)amino]carbonyl]-2-nitrophenyl]amino]methyl]-(2*S*)-1-pyrrolidinecarboxylic acid-1,1-dimethylethyl ester



10


Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (130 mg, 0.373 mmol) and (*S*)-2-aminomethyl-1-Boc-pyrrolidine (82 mg, 0.410 mmol) in 3 mL of EtOH containing Et₃N (0.080 mL, 0.559 mmol). The product was purified by flash chromatography on silica gel using 3:1 / hexanes:EtOAc as eluent. Yield: 165 mg (84%); ¹H NMR (CDCl₃) 1.47 (brs, 9H), 1.83 (m, 1H), 1.90 (m, 2H), 2.03 (m, 1H), 3.31 (m, 1H), 3.37 (m, 2H), 3.62 (m, 1H), 4.09 (m, 2H), 4.24 (m, 6H), 7.26 (m, 1H), 7.50 (m, 1H), 8.27 (s, 1H), 8.55 (brs, 1H); MS (ESI) 529.38 (MH⁺).

15

Intermediate 7

2-[[[4-[[Bis(2,2,2-trifluoroethyl)amino]carbonyl]-2-nitrophenyl]amino]methyl]-(2*R*)-1-pyrrolidinecarboxylic acid-1,1-dimethylethyl ester


20



Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (157 mg, 0.450 mmol) and (*R*)-2-aminomethyl-1-Boc-pyrrolidine (100 mg, 0.495 mmol) in 3 mL of EtOH containing Et₃N (0.095 mL, 0.559 mmol). The product was purified by flash chromatography on silica gel using 3:1 / hexanes:EtOAc as eluent. Yield: 232 mg (98%); ¹H NMR (CDCl₃) 1.47 (brs, 9H), 1.83 (m, 1H), 1.90 (m, 2H), 2.03 (m, 1H), 3.31 (m, 1H), 3.37 (m, 2H), 3.62 (m, 1H), 4.09 (m, 2H), 4.24 (m, 6H), 7.26 (m, 1H), 7.50 (m, 1H), 8.27 (s, 1H), 8.55 (brs, 1H); MS (ESI) 529.38 (MH⁺).

Intermediate 8


3-Nitro-4-[(4-pyridinylmethyl)amino]-*N,N*-bis(2,2,2-trifluoroethyl)-benzamide



4-Fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (120 mg, 0.345 mmol) and 4-(aminomethyl)pyridine (0.070 mL, 0.380 mmol) were stirred in 3 mL of CH₃CN. The solvent was concentrated. The residue was dissolved in EtOAc and washed with saturated NaHCO₃, brine and dried over anhydrous Na₂SO₄. The product was purified by flash chromatography on silica gel using EtOAc as eluent affording the title compound as yellow oil. Yield: 145 mg (79%); ¹H NMR (CDCl₃) 4.19 (q, J = 8.59Hz, 4H), 4.71 (d, J = 6.25Hz, 2H), 6.69 (d, J = 8.79Hz, 1H), 7.45 (m, 3H), 8.32 (s, 1H), 8.65 (d, J = 6.25Hz, 2H), 8.73 (m, 1H); MS (ESI) 437.24 (MH⁺).

Intermediate 9


3-Nitro-4-[[[(tetrahydro-2H-pyran-4-yl)methyl]amino]-N,N-bis(2,2,2-trifluoroethyl)-benzamide



- 5 Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (75 mg, 0.215 mmol) and 4-aminomethyltetrahydropyran (27mg, 0.236 mmol) in 3 mL of EtOH containing Et₃N (0.045 mL, 0.323 mmol). The product was purified by flash chromatography on silica gel using 2:1 / hexanes:EtOAc as eluent. Yield: 87 mg (91%); ¹H NMR (CDCl₃) 1.47 (m, 2H), 1.75 (m, 2H), 1.98 (m, 1H), 3.27 (t, J = 5.47Hz, 2H), 3.43 (m, 2H), 4.03 (m, 2H), 4.24 (q, J = 8.33Hz, 6H), 6.93 (d, J = 8.98Hz, 1H), 7.54 (d, J = 8.98Hz, 1H), 8.30 (s, 1H), 8.40 (brs, 1H); MS (ESI) 444.31 (MH⁺).
- 10

Intermediate 10


- 15 **3-Nitro-4-[[[(2*R*)-tetrahydro-2-furanyl]methyl]amino]-N,N-bis(2,2,2-trifluoroethyl)-benzamide**



- Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (78 mg, 0.224 mmol) and R-(-)-tetrahydrofurfurylamine (0.025 mL, 0.246 mmol) in 3 mL of EtOH containing Et₃N (0.047 mL, 0.336 mmol). The product was purified by flash chromatography on silica gel using 2:1 / hexanes:EtOAc as eluent. Yield: 96 mg (95%); ¹H NMR (CDCl₃) 1.72 (m, 1H), 1.99 (m, 2H), 2.13 (m, 1H), 3.99 (m, 1H), 3.53 (m, 1H), 3.85 (m, 1H), 3.97 (m, 1H), 4.25 (m, 5H), 6.99 (d, J = 8.98Hz, 1H), 7.53 (d, J = 8.98Hz, 1H), 8.30 (s, 1H), 8.51 (brs, 1H); MS (ESI) 430.31 (MH⁺).
- 20
- 25

Intermediate 11

3-Nitro-4-[[[(2*S*)-tetrahydro-2-furanyl]methyl]amino]-*N,N*-bis(2,2,2-trifluoroethyl)-benzamide



5


Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (78 mg, 0.224 mmol) and *S*-(+)-tetrahydrofurfurylamine (0.025 mL, 0.246 mmol) in 3 mL of EtOH containing Et₃N (0.047 mL, 0.336 mmol). The product was purified by flash chromatography on silica gel using 1:1 / hexanes:EtOAc as eluent. Yield: 95 mg (95%); ¹H NMR (CDCl₃) 1.72 (m, 1H), 1.99 (m, 2H), 2.13 (m, 1H), 3.99 (m, 1H), 3.53 (m, 1H), 3.85 (m, 1H), 3.97 (m, 1H), 4.25 (m, 5H), 6.99 (d, *J* = 8.98Hz, 1H), 7.53 (d, *J* = 8.98Hz, 1H), 8.30 (s, 1H), 8.51 (brs, 1H); MS (ESI) 430.28 (MH⁺).

10

15

Intermediate 12

3-Nitro-4-[[[(tetrahydro-2*H*-pyran-2-yl)methyl]amino]-*N,N*-bis(2,2,2-trifluoroethyl)-benzamide




Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (83 mg, 0.238 mmol) and *R/S*-2-aminomethyl-tetrahydropyran hydrochloride (40 mg, 0.262 mmol) in 3 mL of EtOH containing Et₃N (0.070 mL, 0.476 mmol). The product was purified by flash chromatography on silica gel using 2:1 / hexanes:EtOAc as eluent. Yield: 100 mg (95%); ¹H NMR (CDCl₃) 1.48 (m, 1H), 1.58 (m, 3H), 1.68 (m, 1H), 1.95 (m, 1H), 3.35 (m, 1H), 3.41

20

(m, 1H), 3.52 (m, 1H), 3.64 (m, 1H), 4.07 (m, 1H), 4.26 (m, 4H), 6.96 (d, $J = 8.98\text{Hz}$, 1H), 7.54 (d, $J = 8.98\text{Hz}$, 1H), 8.31 (s, 1H), 8.55 (brs, 1H); MS (ESI) 530.21 (MH⁺)

Intermediate 13


- 5 2-[[[4-[[Bis(2,2,2-trifluoroethyl)amino]carbonyl]-2-nitrophenyl]amino]methyl]-(2*R*)-1-piperidinecarboxylic acid-1,1-dimethylethyl ester



- Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (200 mg, 0.574 mmol) and *R*-2-methylamino-1-Boc-piperidine (148 mg, 0.689 mmol) in 5 mL of EtOH containing Et₃N (0.160 mL, 1.14 mmol). The product was purified by flash chromatography on silica gel using 1:1 / hexanes:EtOAc as eluent. Yield: 310 mg (99%); ¹H NMR (CDCl₃) 1.47 (s, 9H), 1.55 (m, 1H), 1.74 (m, 5H), 2.80 (m, 1H), 3.37 (m, 1H), 3.65 (m, 1H), 4.10 (m, 1H), 4.27 (q, $J = 8.40\text{Hz}$, 4H), 4.64 (m, 1H), 7.07 (d, $J = 8.59\text{Hz}$, 1H), 7.55 (d, $J = 8.79\text{Hz}$, 1H), 8.31 (s, 1H), 8.39 (brs, 1H); MS (ESI) 443.35 (MH⁺ - t-Boc).

Intermediate 14


tert-Butyl 3-[[[4-[[bis(2,2,2-trifluoroethyl)amino]carbonyl]-2-nitrophenyl]amino]methyl] morpholine-4-carboxylate



Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (213 mg, 0.612 mmol) and *tert*-butyl 3-(aminomethyl)morpholine-4-carboxylate (160 mg, 0.734 mmol) in 10 mL of EtOH containing TEA (0.130 mL, 0.918 mmol). The product was purified by flash chromatography on silica gel using 2:1 / hexanes:EtOAc as eluent. Yield: 304 mg (91%); ¹H NMR (400 MHz, CHLOROFORM-D) δ 1.49 (s, 9H), 3.17 (m, 1H), 3.52 (m, 1H), 3.64 (m, 3H), 3.92 (m, 2H), 4.25 (q, J = 8.20 Hz, 6H), 7.57 (d, J = 8.59 Hz, 1H), 8.30 (d, J = 1.95 Hz, 1H), 8.48 (m, 1H); MS (ESI) 544.75 (MH⁺).

10 **Intermediate 15**

tert-Butyl (2*S*)-2-[[[4-{[bis(2,2,2-trifluoroethyl)amino]carbonyl}-2-nitrophenyl)amino]methyl] piperidine-1-carboxylate




15

Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (200 mg, 0.574 mmol) and *tert*-butyl (2*S*)-2-(aminomethyl)piperidine-1-carboxylate (150 mg, 0.689 mmol) in 10 mL of EtOH containing TEA (0.120 mL, 0.861 mmol). The product was purified by flash chromatography on silica gel using 3:1 / hexanes:EtOAc as eluent. Yield: 311 mg (99%); ¹H NMR (400 MHz, CHLOROFORM-D) δ 1.47 (s, 9H), 1.52 (m, 1H), 1.66 (m, 1H), 1.71 (s, 2H), 1.74 (s, 2H), 2.79 (m, 1H), 3.36 (m, 1H), 3.65 (m, 1H), 4.09 (s, 1H), 4.26 (q, J = 8.33 Hz, 4H), 4.64 (m, 1H), 7.07 (d, J = 8.98 Hz, 1H), 7.55 (dd, J = 8.89, 2.05 Hz, 1H), 8.30 (d, J = 2.15 Hz, 1H), 8.39 (m, 1H); MS (ESI) 542.81 (MH⁺).

25

Intermediate 16


4-[(Cyclobutylmethyl)amino]-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide



Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (237 mg, 0.680 mmol) and cyclobutylmethyl amine (0.205 mL of a 4M/MeOH solution, 0.816 mmol) in 3 mL of EtOH containing TEA (0.140 mL, 1.02 mmol). The product was purified by flash chromatography on silica gel using 3:1 / hexanes:EtOAc as eluent. Yield: 262 mg (93%); ¹H NMR (400 MHz, CHLOROFORM-D) δ 1.83 (m, 2H), 1.97 (m, 1H), 2.02 (m, 1H), 2.20 (m, 2H), 2.73 (m, 1H), 3.37 (m, 2H), 4.27 (m, 4H), 6.93 (d, J = 8.98 Hz, 1H), 7.53 (d, J = 8.79 Hz, 1H), 8.24 (m, 1H), 8.30 (m, 1H); MS (ESI) 413.95 (MH⁺).

Intermediate 17

4-[(Cyclopentylmethyl)amino]-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide




Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (122 mg, 0.350 mmol) and cyclopentylmethyl amine (42 mg, 0.420 mmol) in 3 mL of EtOH containing TEA (0.075 mL, 0.525 mmol). The product was purified by flash chromatography on silica gel using 3:1 / hexanes:EtOAc as eluent. Yield: 141 mg (94%); ¹H NMR (400 MHz, CHLOROFORM-D) δ 1.30 (m, 2H), 1.64 (m, 4H), 1.89 (m, 2H), 2.26 (m, 1H), 3.24 (dd, J = 7.23, 5.04 Hz, 2H), 4.23 (q, J = 8.42 Hz, 4H), 6.91 (d, J = 8.97 Hz, 1H), 7.50

(dd, $J = 8.97, 2.20$ Hz, 1H), 8.26 (d, $J = 2.20$ Hz, 1H), 8.33 (s, 1H); MS (ESI) 427.82 (MH⁺).

Intermediate 18


5 4-[(3-Furylmethyl)amino]-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide



Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (103 mg, 0.296 mmol) and 3-furylmethylamine (35 mg, 0.355 mmol) in 3 mL of EtOH containing TEA (0.060 mL, 0.444 mmol). The product was purified by flash chromatography on silica gel using 3:1 / hexanes:EtOAc as eluent. Yield: 114 mg (91%); ¹H NMR (400 MHz, CHLOROFORM-D) δ 4.26 (q, $J = 8.40$ Hz, 4H), 4.44 (d, $J = 5.27$ Hz, 2H), 6.43 (m, 1H), 7.00 (d, $J = 8.98$ Hz, 1H), 7.46 (m, 2H), 7.55 (dd, $J = 8.88, 2.05$ Hz, 1H), 8.32 (d, $J = 2.15$ Hz, 1H), 8.45 (m, 1H); MS (ESI) 425.72 (MH⁺).

Intermediate 19

20 3-Nitro-4-[(3-thienylmethyl)amino]-*N,N*-bis(2,2,2-trifluoroethyl)benzamide




Following the general procedure for **Intermediate 2** using 4-fluoro-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (125 mg, 0.359 mmol) and 3-thienylmethylamine (49 mg, 0.431 mmol) in 3 mL of EtOH containing TEA (0.075 mL, 0.539 mmol).

The product was purified by flash chromatography on silica gel using 3:1 /
hexanes:EtOAc as eluent. Yield: 143 mg (90%); ¹H NMR (400 MHz,
CHLOROFORM-D) δ 4.24 (q, J = 8.49 Hz, 4H), 4.59 (d, J = 5.13 Hz, 2H), 6.95 (d, J
= 8.79 Hz, 1H), 7.07 (dd, J = 5.04, 1.37Hz, 1H), 7.22 (dd, J = 3.02, 1.19 Hz, 1H), 7.37
5 (dd, J = 5.04, 3.02 Hz, 1H), 7.51 (dd, J = 8.61, 1.83 Hz, 1H), 8.31 (d, J = 2.01 Hz,
1H), 8.60 (m, 1H); MS (ESI) 441.75 (MH⁺).

Intermediate 20


4-[[[(2*R*)-1-Methyl-2-pyrrolidinyl]methyl]amino]-3-nitro-*N,N*-bis(2,2,2-
10 trifluoroethyl)-benzamide



Intermediate 7 (167 mg, 0.308 mmol) was stirred in 2 mL of 1M HCl/AcOH at RT
15 for 1h. The solvent was evaporated. The residue was dissolved in THF (5 mL) and
an excess of 37% HCHO/H₂O (1mL) was added followed by NaBH(OAc)₃ (130 mg,
0.616 mmol). The solution was stirred at RT for 1h. The solvent was concentrated.
The residue was dissolved in EtOAc and washed with saturated NaHCO₃, brine and
dried over anhydrous Na₂SO₄. The solvent was concentrated and the product was
20 dried under vacuum to give the titled compound. Yield: 136 mg (99%); ¹H NMR
(CDCl₃) δ 1.69 (m, 2H), 1.81 (m, 1H), 2.01 (m, 1H), 2.28 (m, 1H), 2.35 (s, 3H), 2.58
(m, 1H), 3.15 (m, 1H), 3.32 (m, 2H), 4.21 (q, J = 8.49Hz, 4H), 6.88 (d, J = 8.97Hz,
1H), 7.49 (d, J = 8.88Hz, 1H), 8.26 (s, 1H), 8.61 (brs, 1H). MS (ESI) 443.95 (MH⁺).

Intermediate 21


3-Amino-4-[(3-methylbutyl)amino]-*N,N*-bis(2,2,2-trifluoroethyl)-benzamide



- 4-[(3-Methylbutyl)amino]-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)-benzamide (190 mg, 0.457 mmol) was dissolved in EtOAc (10 mL) containing a catalytic amount of 10% Pd/C. The solution was shaken with a Parr hydrogenation apparatus under H₂ atmosphere (40 psi) at RT overnight. The solution was filtered through diatomaceous earth and the solvent was concentrated giving the intermediate 21 as a white foam (176 mg, 99%); MS (ESI) 386.17 (MH⁺).

Intermediate 22

- 10 **3-Amino-4-[[[(2*R*)-1-methyl-2-pyrrolidiny]methyl]amino]-*N,N*-bis(2,2,2-trifluoroethyl)-benzamide**



- 15 Following the same procedure as for **Intermediate 21** using intermediate 20 (130 mg, 0.294 mmol) as starting material yielded the titled intermediate 16. Yield: 101 mg (85%); MS (ESI) 413.18 (MH⁺).

- 20 The synthesis of all other intermediates followed the same hydrogenation procedure as for **Intermediate 21** giving the desired products in quantitative yields (to be included in Table 2).











Table 2. Intermediates prepared following a general procedure for Intermediate 21:

<u>Intermediates</u>	<u>R</u>	<u>(MH⁺)</u>
<u>Intermediate 23</u>		370.17
<u>Intermediate 24</u>		412.36
<u>Intermediate 25</u>		396.25
<u>Intermediate 26</u>		499.44
<u>Intermediate 27</u>		499.44
<u>Intermediate 28</u>		407.32
<u>Intermediate 29</u>		414.36
<u>Intermediate 30</u>		400.17
<u>Intermediate 31</u>		400.17

<u>Intermediate 32</u>		414.36
<u>Intermediate 33</u>		513.45
<u>Intermediate 34</u>		515.11
<u>Intermediate 35</u>		512.92
<u>Intermediate 36</u>		384.15
<u>Intermediate 37</u>		397.92

Intermediate 38


3-amino-4-[(3-furylmethyl)amino]-*N,N*-bis(2,2,2-trifluoroethyl)benzamide



5

4-[(3-Furylmethyl)amino]-3-nitro-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (108 mg, 0.254 mmol) was dissolved in 5 mL of DMF. Tin (II) chloride dihydrate (860 mg, 1.27 mmol) was added and the solution was stirred under nitrogen at RT for 24h. The solvent was evaporated and the residue was taken up in EtOAc. The organic phase was washed with saturated NaHCO₃ solution, brine and dried over anhydrous MgSO₄. The crude product was used directly for the next step. Yield: 80 mg (80%); MS (ESI) 396.11 (MH⁺).


10

Intermediate 39**3-Amino-4-[(3-thienylmethyl)amino]-*N,N*-bis(2,2,2-trifluoroethyl)benzamide**

5

Same procedure used as above using 3-nitro-4-[(3-thienylmethyl)amino]-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (138 mg, 0.312 mmol) and tin (II) chloride dihydrate (210 mg, 0.936 mmol) in 5 mL of DMF. Yield: 125 mg (97%); MS (ESI) 412.08 (MH⁺).

10

Example 3**2-[(4-Ethoxyphenyl)methyl]-1-(3-methylbutyl)-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide**

15


3-Amino-4-[(3-methylbutyl)amino]-*N,N*-bis(2,2,2-trifluoroethyl)-benzamide (176 mg, 0.457 mmol), 4-ethoxyphenylacetyl chloride (90 mg, 0.457 mmol) and zinc dust (30 mg, 0.457 mmol) were stirred in 1,2-dichloroethane (3 mL) at RT for 30 min. A catalytic amount of concentrated HCl (11.6 M) was added and the solution was stirred at 85 °C overnight. The solution was cooled to RT and diluted with dichloromethane. The organic phase was washed with saturated NaHCO₃ aqueous solution, brine and dried over anhydrous Na₂SO₄. The product was purified by reversed-phase HPLC using 20-80% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 172 mg (71%); ¹H NMR (CD₃OD) 0.90 (d, J =

20

6.64Hz, 6H), 1.35 (t, J = 7.03Hz, 3H), 1.61 (m, 1H), 4.00 (q, J = 7.03Hz, 2H), 4.23 (m, 2H), 4.39 (m, 6H), 6.90 (d, J = 8.79Hz, 2H), 7.20 (d, J = 8.79Hz, 2H), 7.44 (d, J = 8.49Hz, 1H), 7.66 (d, J = 8.40Hz, 1H), 7.75 (s, 1H); MS (ESI) 530.21 (MH⁺); Anal. Calcd. for C₂₆H₂₉N₃O₂F₆ + 0.3 TFA + 0.2 H₂O: C, 56.31; H, 5.28; N, 7.41. Found: C, 56.29; H, 5.12; N, 7.48.

Example 4


1-(Cyclopropylmethyl)-2-[(4-ethoxyphenyl)methyl]-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide



Following the general procedure in **Example 3** using intermediate 23 (77 mg, 0.210 mmol), 4-ethoxyphenylacetyl chloride (42 mg, 0.210 mmol) and zinc dust (14 mg, 0.210 mmol) in 3 mL of 1,2-dichloroethane. The product was purified by reversed-phase HPLC using 20-80% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 74 mg (73%); ¹H NMR (CD₃OD) 0.47 (m, 2H), 0.63 (m, 2H), 1.27 (m, 1H), 1.38 (t, J = 6.93Hz, 3H), 4.02 (q, J = 6.90Hz, 2H), 4.37 (m, 6H), 4.56 (s, 2H), 6.97 (d, J = 8.59Hz, 2H), 7.28 (d, J = 8.59Hz, 2H), 7.60 (d, J = 8.50Hz, 1H), 7.79 (s, 1H), 8.00 (d, J = 8.40Hz, 1H); MS (ESI) 514.22 (MH⁺); Anal. Calcd. for C₂₅H₂₅N₃O₂F₆ + 1.0 TFA + 0.1 H₂O: C, 51.53; H, 4.20; N, 6.68; Found: C, 51.49; H, 4.18; N, 6.55.

Example 5


1-(Cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide



Following the general procedure in **Example 3** using intermediate 24 (95 mg, 0.227 mmol), 4-ethoxyphenylacetyl chloride (50 mg, 0.250 mmol) and zinc dust (16 mg, 0.250 mmol) in 3 mL of 1,2-dichloroethane. The product was purified by reversed-phase HPLC using 20-80% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 72mg (48%); ¹H NMR (CD₃OD) 1.12 (m, 5H), 1.36 (t, J = 7.03Hz, 3H), 1.58 (m, 2H), 1.69 (m, 4H), 4.02 (q, J = 7.03Hz, 2H), 4.23 (d, J = 7.62Hz, 2H), 4.36 (m, 4H), 4.50 (s, 2H), 6.94 (d, J = 8.79Hz, 2H), 7.24 (d, J = 8.79Hz, 2H), 7.56 (d, J = 8.59Hz, 1H), 7.76 (s, 1H), 7.90 (d, J = 8.59Hz, 1H); MS (ESI) 556.47 (MH⁺); Anal. Calcd. for C₂₈H₃₁N₃O₂F₆ + 0.9 TFA + 0.2 H₂O; C, 54.09; H, 4.92; N, 6.35.; Found: C, 54.12; H, 4.74; N, 6.20.

Example 6

2-[(4-Ethoxyphenyl)methyl]-1-(2-furanylmethyl)-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide




Following the general procedure in **Example 3** using intermediate 25 (106 mg, 0.270 mmol), 4-ethoxyphenylacetyl chloride (59 mg, 0.297 mmol) and zinc dust (19 mg, 0.297 mmol) in 3 mL of 1,2-dichloroethane. The product was purified by reversed-phase HPLC using 20-80% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 78 mg (45%); ¹H NMR (CD₃OD) 1.34 (t, J = 7.03Hz, 3H), 3.98 (q, J = 7.03Hz, 2H), 4.36 (m, 4H), 4.46 (s, 2H), 5.48 (s, 2H), 6.31 (m, 2H), 6.86 (d, J = 8.79Hz, 2H), 7.17 (d, J = 8.79Hz, 2H), 7.41 (m, 2H),

7.70 (s, 1H), 7.79 (d, $J = 8.59\text{Hz}$, 1H); MS (ESI) 540.38 (MH⁺); Anal. Calcd. for $\text{C}_{26}\text{H}_{23}\text{N}_3\text{O}_3\text{F}_6 + 0.4 \text{ TFA}$; C, 55.02; H, 4.03; N, 7.15. Found: C, 55.22; H, 4.21; N, 6.75.

5 Example 7


2-[(4-Ethoxyphenyl)methyl]-1-[(2*S*)-2-pyrrolidinylmethyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide



Following the general procedure in **Example 3** using intermediate 27 (210 mg, 0.420 mmol), 4-ethoxyphenylacetyl chloride (85 mg, 0.420 mmol) and zinc dust (27 mg, 0.420 mmol) in 3 mL of 1,2-dichloroethane. The product was purified by reversed-phase HPLC using 10-50% $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 95 mg (35%); ^1H NMR (CD_3OD) 1.35 (t, $J = 7.03\text{Hz}$, 3H), 1.78 (m, 1H), 1.99 (m, 1H), 2.13 (m, 2H), 3.22 (m, 1H), 3.44 (m, 1H), 3.79 (m, 1H), 4.00 (q, $J = 7.03\text{Hz}$, 2H), 4.36 (m, 4H), 4.43 (s, 2H), 4.64 (d, $J = 6.84\text{Hz}$, 2H), 6.91 (d, $J = 8.79\text{Hz}$, 2H), 7.22 (d, $J = 8.79\text{Hz}$, 2H), 7.49 (d, $J = 8.49\text{Hz}$, 1H), 7.76 (m, 2H); MS (ESI) 543.44 (MH⁺); Anal. Calcd. for $\text{C}_{26}\text{H}_{28}\text{N}_4\text{O}_2\text{F}_6 + 2.2 \text{ TFA} + 0.3 \text{ H}_2\text{O}$; C, 45.71; H, 3.89; N, 7.01. Found: C, 45.64; H, 3.74; N, 7.30.

20 Example 8

2-[(4-Ethoxyphenyl)methyl]-1-[(2*R*)-2-pyrrolidinylmethyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide




Following the general procedure in **Example 3** using intermediate 26 (150 mg, 0.301 mmol), 4-ethoxyphenylacetyl chloride (60 mg, 0.301 mmol) and zinc dust (20 mg, 0.301 mmol) in 3 mL of 1,2-dichloroethane. The product was purified by reversed-

- 5 compound as the corresponding TFA salt. Yield: 83 mg (42%); ^1H NMR (CD_3OD) 1.35 (t, $J = 7.03\text{Hz}$, 3H), 1.78 (m, 1H), 1.99 (m, 1H), 2.13 (m, 2H), 3.22 (m, 1H), 3.44 (m, 1H), 3.79 (m, 1H), 4.00 (q, $J = 7.03\text{Hz}$, 2H), 4.36 (m, 4H), 4.43 (s, 2H), 4.64 (d, $J = 6.84\text{Hz}$, 2H), 6.91 (d, $J = 8.79\text{Hz}$, 2H), 7.22 (d, $J = 8.79\text{Hz}$, 2H), 7.49 (d, $J = 8.49\text{Hz}$, 1H), 7.76 (s, 1H), 7.81 (d, $J = 8.40\text{Hz}$, 1H); MS (ESI) 543.44 (MH^+); Anal. Calcd. for $\text{C}_{26}\text{H}_{28}\text{N}_4\text{O}_2\text{F}_6 + 2.2 \text{ TFA} + 0.4 \text{ H}_2\text{O}$: C, 45.61; H, 3.90; N, 7.00. Found: C, 45.57; H, 3.74; N, 7.30.

Example 9


- 2-[(4-Ethoxyphenyl)methyl]-1-(4-pyridinylmethyl)-*N,N*-bis(2,2,2-trifluoroethyl)-
15 1*H*-benzimidazole-5-carboxamide



- Following the general procedure in **Example 3** using intermediate 28 (130 mg, 0.320 mmol), 4-ethoxyphenylacetyl chloride (70 mg, 0.352 mmol) and zinc dust (23 mg, 0.352 mmol) in 3 mL of 1,2-dichloroethane. The product was purified by reversed-
20 phase HPLC using 10-50% $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 30 mg (15%); ^1H NMR (CD_3OD) 1.30 (t, $J = 7.03\text{Hz}$, 3H), 3.88 (q, $J = 7.03\text{Hz}$, 2H), 4.36 (m, 4H), 4.41 (s, 2H), 4.64 (d, $J = 6.84\text{Hz}$, 2H), 5.91 (s, 2H), 6.62 (d, $J = 8.40\text{Hz}$, 2H), 7.05 (d, $J = 8.59\text{Hz}$, 2H), 7.28 (d, $J = 5.66\text{Hz}$, 2H), 7.43 (d, $J = 8.40\text{Hz}$, 1H), 7.60 (d, $J = 8.40\text{Hz}$, 1H), 7.86 (s, 1H),
25 8.52 (brs, 2H); MS (ESI) 551.43 (MH^+); Anal. Calcd. for $\text{C}_{27}\text{H}_{24}\text{N}_4\text{O}_2\text{F}_6 + 2.5 \text{ TFA}$; C, 46.00; H, 3.20; N, 6.71. Found: C, 46.17; H, 3.11; N, 6.63.

Example 10


2-[1-(4-Ethoxyphenyl)ethyl]-1-(4-pyridinylmethyl)-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide



Following the general procedure in **Example 3** using intermediate 28 (78 mg, 0.193 mmol), 4-ethoxy- \square -methyl-phenylacetyl chloride (45 mg, 0.231 mmol) and zinc dust (15 mg, 0.231 mmol) in 3 mL of 1,2-dichloroethane. The product was purified by reversed-phase HPLC using 10-50% $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 16mg (15%). ^1H NMR (CD_3OD) 1.29 (t, $J = 6.93\text{Hz}$, 3H), 1.77 (d, $J = 7.03\text{Hz}$, 3H), 3.85 (q, $J = 7.03\text{Hz}$, 2H), 4.37 (m, 5H), 4.50 (m, 1H), 5.67 (d, $J = 18.75\text{Hz}$, 1H), 5.82 (d, $J = 18.75\text{Hz}$, 1H), 6.56 (d, $J = 8.79\text{Hz}$, 2H), 6.96 (d, $J = 8.79\text{Hz}$, 2H), 7.05 (d, $J = 6.25\text{Hz}$, 1H), 7.34 (d, $J = 8.40\text{Hz}$, 1H), 7.46 (d, $J = 8.40\text{Hz}$, 1H), 7.86 (s, 1H), 8.40 (brs, 2H); MS (ESI) 565.43 (MH^+); Anal. Calcd. for $\text{C}_{28}\text{H}_{26}\text{N}_4\text{O}_2\text{F}_6 + 1.6 \text{ TFA}$; C, 50.17; H, 3.72; N, 7.50. Found: C, 50.20; H, 3.71; N, 7.44.

Example 11

2-[(4-Ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide




Following the general procedure in **Example 3** using intermediate 29 (75 mg, 0.181 mmol), 4-ethoxyphenylacetyl chloride (40 mg, 0.199 mmol) and zinc dust (13 mg, 0.199 mmol) in 3 mL of 1,2-dichloroethane. The product was purified by reversed-phase HPLC using 10-60% $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ and then lyophilized affording the title

compound as the corresponding TFA salt. Yield: 70 mg (58%). ^1H NMR (CD_3OD) 1.38 (t, $J = 7.03\text{Hz}$, 3H), 1.47 (m, 5H), 2.02 (m, 2H), 3.24 (m, 3H), 3.90 (m, 2H), 4.03 (q, $J = 7.03\text{Hz}$, 2H), 4.33 (d, $J = 7.62\text{Hz}$, 2H), 4.38 (m, 4H), 4.54 (s, 2H), 6.96 (d, $J = 8.59\text{Hz}$, 2H), 7.28 (d, $J = 8.59\text{Hz}$, 2H), 7.58 (d, $J = 8.49\text{Hz}$, 1H), 7.78 (s, 1H), 7.97 (d, $J = 8.40\text{Hz}$, 1H); MS (ESI) 558.48 (MH^+); Anal. Calcd. for $\text{C}_{27}\text{H}_{29}\text{N}_3\text{O}_3\text{F}_6 + 1.2 \text{ TFA} + 0.2 \text{ H}_2\text{O}$; C, 50.59; H, 4.42; N, 6.02. Found: C, 50.54; H, 4.47; N, 6.00.

Example 12


2-[(4-Ethoxyphenyl)methyl]-1-[(2*R*)-tetrahydro-2-furanyl]methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide



Following the general procedure in Example 3 using intermediate 30 (88 mg, 0.220 mmol), 4-ethoxyphenylacetyl chloride (48 mg, 0.242 mmol) and zinc dust (16 mg, 0.242 mmol) in 3 mL of 1,2-dichloroethane. The product was purified by reversed-phase HPLC using 10-60% $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 70 mg (50%). ^1H NMR (CD_3OD) 1.38 (t, $J = 7.03\text{Hz}$, 3H), 1.74 (m, 1H), 1.97 (m, 2H), 2.18 (m, 1H), 3.72 (q, $J = 7.23\text{Hz}$, 1H), 3.91 (q, $J = 7.42\text{Hz}$, 1H), 4.03 (q, $J = 7.03\text{Hz}$, 2H), 4.21 (m, 1H), 4.36 (br s, 4H), 4.52 (m, 1H), 4.60 (s, 2H), 4.67 (m, 1H), 6.97 (d, $J = 8.40\text{Hz}$, 2H), 7.29 (d, $J = 8.40\text{Hz}$, 2H), 7.61 (d, $J = 8.59\text{Hz}$, 1H), 7.76 (s, 1H), 8.03 (d, $J = 8.59\text{Hz}$, 1H); MS (ESI) 544.45 (MH^+); Anal. Calcd. for $\text{C}_{26}\text{H}_{27}\text{N}_3\text{O}_3\text{F}_6 + 1.4 \text{ TFA} + 0.2 \text{ H}_2\text{O}$; C, 48.95; H, 4.11; N, 5.95. Found: C, 48.95; H, 3.93; N, 6.00.

Example 13


2-[(4-Ethoxyphenyl)methyl]-1-[(2*S*)-tetrahydro-2-furanyl]methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide



Following the general procedure in **Example 3** using intermediate 31 (85 mg, 0.212 mmol), 4-ethoxyphenylacetyl chloride (46 mg, 0.233 mmol) and zinc dust (15 mg, 0.233 mmol) in 3 mL of 1,2-dichloroethane. The product was purified by reversed-phase HPLC using 10-60% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 70mg (50%). ¹H NMR (CD₃OD) 1.38 (t, J = 7.03Hz, 3H), 1.74 (m, 1H), 1.95 (m, 2H), 2.18 (m, 1H), 3.72 (q, J = 7.23Hz, 1H), 3.91 (q, J = 7.42Hz, 1H), 4.03 (q, J = 7.03Hz, 2H), 4.21 (m, 1H), 4.38 (brs, 4H), 4.49 (m, 1H), 4.60 (s, 2H), 4.66 (d, 1H), 6.98 (d, J = 8.40Hz, 2H), 7.29 (d, J = 8.40Hz, 2H), 7.61 (d, J = 8.59Hz, 1H), 7.76(s, 1H), 8.02 (d, J = 8.59Hz, 1H); MS (ESI) 544.45 (MH⁺); Anal. Calcd. for C₂₆H₂₇N₃O₃F₆ + 1.4 TFA + 0.1 H₂O; C, 49.07; H, 4.09; N, 5.96. Found: C, 49.08; H, 4.05; N, 6.11.

Example 14

2-[(4-Ethoxyphenyl)methyl]-1-[(tetrahydro-2H-pyran-2-yl)methyl]-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide




Following the general procedure in **Example 3** using intermediate 32 (93 mg, 0.225 mmol), 4-ethoxyphenylacetyl chloride (49 mg, 0.248 mmol) and zinc dust (16 mg, 0.248 mmol) in 3 mL of 1,2-dichloroethane. The product was purified by reversed-phase HPLC using 10-60% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 58 mg (38%). ¹H NMR (CD₃OD) 1.38 (t, J = 7.03Hz, 3H), 1.42 (m, 1H), 1.51 (m, 3H), 1.80 (m, 1H), 1.90 (m, 1H), 3.23

(m, 1H), 3.59 (m, 1H), 3.89 (m, 1H), 4.04 (q, J = 7.03Hz, 2H), 4.30 (brs, 4H), 4.48 (m, 1H), 4.5 (s, 1H), 4.60 (d, J = 8.59Hz, 2H), 6.97 (d, J = 8.79Hz, 2H), 7.27 (d, J = 8.59Hz, 2H), 7.61 (d, J = 8.59Hz, 1H), 7.76 (s, 1H), 8.01 (d, J = 8.59Hz, 1H); MS (ESI) 558.53 (MH⁺); Anal. Calcd. for C₂₇H₂₉N₃O₃F₆ + 1.6 TFA; C, 49.02; H, 4.17; N, 5.68. Found: C, 49.12; H, 4.05; N, 5.81.

Example 15


2-[(4-Ethoxyphenyl)methyl]-1-[(2R)-2-piperidinylmethyl]-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide



Following the general procedure in **Example 3** using intermediate 33 (261 mg, 0.509 mmol), 4-ethoxyphenylacetyl chloride (101 mg, 0.509 mmol) and zinc dust (33 mg, 0.509 mmol) in 5 mL of 1,2-dichloroethane. The product was purified by reversed-phase HPLC using 10-60% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 121mg (35%). ¹H NMR (CD₃OD) 1.37 (t, J = 7.03Hz, 3H), 1.45 (m, 1H), 1.58 (m, 1H), 1.65 (m, 1H), 1.88 (brd, 3H), 2.80 (m, 1H), 3.38 (m, 2H), 4.02 (q, J = 7.03Hz, 2H), 4.40 (m, 4H), 4.46 (s, 2H), 4.53 (m, 2H), 6.95 (d, J = 8.79Hz, 2H), 7.24 (d, J = 8.79Hz, 2H), 7.51 (d, J = 8.40Hz, 1H), 7.77 (s, 1H), 7.82 (d, J = 8.40Hz, 1H); MS (ESI) 557.47 (MH⁺); Anal. Calcd. for C₂₇H₃₀N₄O₂F₆ + 2.1 TFA + 0.2 H₂O; C, 46.87; H, 4.10; N, 7.01. Found: C, 46.80; H, 3.90; N, 7.18.

Example 16

2-[(5-Ethoxy-2-pyridyl)methyl]-1-[(tetrahydro-2H-pyran-4-yl)methyl]-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide




3-Amino-4-[[[(tetrahydro-2H-pyran-4-yl)methyl]amino]-N,N-bis(2,2,2-trifluoroethyl)-benzamide (150mg, 0.363 mmol), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (151 mg, 0.399 mmol) and 5-ethoxy-2-pyridylacetic acid (72 mg, 0.399 mmol) were stirred in DMF (5 mL) containing diisopropylethylamine (DIPEA) (0.095 mL, 0.545 mmol) at RT for 3h. The solvent was concentrated and the residue dissolved in EtOAc. The organic phase was washed with saturated aqueous NaHCO₃ solution, brine and dried over anhydrous Na₂SO₄. The solvent was concentrated and the product dissolved in 1,2-dichloroethane (3 mL).

A catalytic amount of concentrated HCl (11.6M) was added and the solution was stirred at 85 °C for 3h. The solvent was concentrated and the residue dissolved in EtOAc. The organic phase was washed with saturated NaHCO₃ solution, brine and dried over anhydrous Na₂SO₄. The product was purified by reversed-phase HPLC using 10-50% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 131 mg (54%). ¹H NMR (CD₃OD) 1.42 (t, J = 7.03 Hz, 3H), 1.51 (m, 5H), 2.14 (m, 1H), 3.25 (m, 1H), 3.90 (d, J = 10.94 Hz, 2H), 4.14 (q, J = 7.03 Hz, 2H), 4.40 (m, 7H), 4.76 (m, 1H), 7.57 (m, 3H), 7.80 (s, 1H), 8.01 (d, J = 8.59 Hz, 1H), 8.24 (s, 1H); MS (ESI) 559.48 (MH⁺); Anal. Calcd. for C₂₆H₂₈N₄O₃F₆ + 1.6 TFA; C, 47.33; H, 4.03; N, 7.56. Found: C, 47.31; H, 4.08; N, 7.60.

Example 17

2-[(5-Ethoxy-2-pyridyl)methyl]-1-(3-methylbutyl)-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide



Following the general procedure in **Example 16** using intermediate 21 (180 mg, 0.469 mmol), O-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyl uronium


hexafluorophosphate (HATU) (196 mg, 0.516 mmol) and 5-ethoxy-2-pyridylacetic acid (93 mg, 0.516 mmol) were stirred in DMF (5 mL) containing

diisopropylethylamine (DIPEA) (0.125 mL, 0.704 mmol). The product was purified by reversed-phase HPLC using 10-50% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 208 mg (69%). ¹H NMR

(CD₃OD) 0.97 (d, *J* = 6.44 Hz, 6H), 1.40 (t, *J* = 7.03 Hz, 3H), 1.58 (m, 2H), 1.70 (m, 1H), 4.12 (q, *J* = 7.03 Hz, 2H), 4.42 (m, 8H), 7.47 (m, 1H), 7.51 (d, *J* = 8.59 Hz, 1H), 7.58 (d, *J* = 8.59 Hz, 1H), 7.81 (s, 1H), 7.88 (d, *J* = 8.59 Hz, 1H), 8.20 (s, 1H); MS (ESI) 531.48 (MH⁺); Anal. Calcd. for C₂₅H₂₈N₄O₂F₆ + 0.8 TFA + 0.1 H₂O; C, 51.24; H, 4.69; N, 8.99. Found: C, 51.30; H, 4.63; N, 8.90.

Example 18

2-[(4-Ethoxyphenyl)methyl]-1-[[2-(2,2,2-trifluoroethyl)amino]methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide




2-[[[2-Amino-4-[[bis(2,2,2-trifluoroethyl)amino]carbonyl]phenyl]amino]methyl]-

(2*R*)-1-pyrrolidinecarboxylic acid-1,1-dimethylethyl ester (80 mg, 0.161 mmol), 4-ethoxyphenylacetyl chloride (32 mg, 0.161 mmol) and zinc dust (11 mg, 0.161 mmol) were stirred in 1,2-dichloroethane (3 mL) at RT for 30 min. A catalytic amount of concentrated HCl (11.6 M) was added and the solution was stirred at 85 °C overnight.

The solution was diluted with DCM and washed with saturated NaHCO₃ solution, brine and dried over anhydrous Na₂SO₄. The solvent was evaporated. The residue was dissolved in MeOH (3 mL) containing a few drops of acetic acid and 37% HCHO/H₂O (1 mL, excess) followed by NaCNBH₃ (12 mg, 0.193 mmol). The solution was stirred at RT for 1h. The solvent was evaporated and the residue was dissolved in EtOAc. The organic phase was washed with saturated NaHCO₃ aqueous solution, brine and dried over anhydrous Na₂SO₄. The product was purified by reversed-phase HPLC using 10-50% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 21 mg (20%); ¹H NMR (CD₃OD) 1.36 (t, J = 7.03 Hz, 3H), 1.82 (m, 1H), 2.01 (m, 2H), 2.94 (s, 3H), 3.18 (m, 1H), 3.72 (m, 1H), 3.79 (m, 1H), 4.00 (q, J = 7.03 Hz, 2H), 4.38 (m, 4H), 4.43 (s, 2H), 4.57 (m, 1H), 4.79 (m, 1H), 6.92 (d, J = 8.79 Hz, 2H), 7.21 (d, J = 8.79 Hz, 2H), 7.46 (d, J = 8.49 Hz, 1H), 7.76 (m, 2H); MS (ESI) 557.50 (MH⁺); Anal. Calcd. for C₂₇H₃₀N₄O₂F₆ + 3.0 TFA + 0.9 H₂O; C, 43.33; H, 3.83; N, 6.12. Found: C, 43.33; H, 3.75; N, 6.20.

Example 19

2-[(4-Ethoxyphenyl)methyl]-1-[(2R)-1-methyl-2-piperidiny]methyl]-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide




2-[(4-Ethoxyphenyl)methyl]-1-[(2R)-2-piperidiny]methyl]-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide (TFA salt) (50 mg, 0.637 mmol) was dissolved in 5mL of THF containing a few drops of glacial acetic acid and an excess of 37% HCHO/H₂O (1 mL). NaBH(OAc)₃ (27 mg, 1.27 mmol) was added and the solution was stirred at RT for 1h. The solvent was evaporated and the residue was dissolved in EtOAc. The organic phase was washed with saturated NaHCO₃ aqueous solution, brine and dried over anhydrous Na₂SO₄. The product was purified by reversed-phase HPLC using 10-50% CH₃CN/H₂O and then lyophilized affording the

title compound as the corresponding TFA salt. Yield: 32 mg (74%); ^1H NMR (CD₃OD) 1.20 (m, 1H), 1.36 (t, J = 7.03 Hz, 3H), 1.44 (m, 1H), 1.72 (m, 2H), 1.86 (m, 1H), 2.95 (m, 1H), 3.03 (s, 3H), 3.24 (m, 1H), 3.57 (m, 1H), 4.01 (q, J = 7.03 Hz, 2H), 4.39 (m, 7H), 6.92 (d, J = 8.59 Hz, 2H), 7.21 (d, J = 8.59 Hz, 2H), 7.44 (d, J = 8.40 Hz, 1H), 7.68 (d, J = 8.20 Hz, 1H), 7.76 (s, 1H); MS (ESI) 571.55 (MH⁺); Anal. Calcd. for C₂₈H₃₂N₄O₂F₆ + 1.5 TFA + 0.2 H₂O; C, 49.96; H, 4.59; N, 7.52. Found: C, 49.97; H, 4.55; N, 7.59.

Example 20

- 10 **2-[(5-Ethoxy-2-pyridyl)methyl]-1-[(2R)-2-pyrrolidinylmethyl]-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide**




Following the general procedure in **Example 16** using intermediate 26 (145 mg, 0.291 mmol), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyl uranium

- 15 hexafluorophosphate (HATU) (125 mg, 0.320 mmol) and 5-ethoxy-2-pyridylacetic acid (60 mg, 0.320 mmol) were stirred in DMF (5 mL) containing diisopropylethylamine (DIPEA) (0.085 mL, 0.495 mmol). The product was purified by reversed-phase HPLC using 10-50% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 96 mg (43%). ^1H NMR (CD₃OD) 1.40 (t, J = 7.03 Hz, 3H), 1.99 (m, 1H), 2.12 (m, 1H), 2.23 (m, 1H), 2.38 (m, 1H), 3.39 (m, 1H), 3.52 (m, 1H), 4.12 (q, J = 7.03 Hz, 2H), 4.36 (m, 6H), 4.82 (m, 1H), 4.92 (m, 1H), 7.48 (d, J = 8.59 Hz, 1H), 7.58 (m, 2H), 7.71 (s, 1H), 7.88 (d, J = 8.40 Hz, 1H), 8.17 (s, 1H); MS (ESI) 544.45 (MH⁺); Anal. Calcd. for C₂₅H₂₇N₅O₂F₆ + 2.2 TFA + 0.2 H₂O; C, 44.25; H, 3.74; N, 8.78. Found: C, 44.22; H, 3.77; N, 8.78.

25

Example 21


- 2-[1-(4-Ethoxyphenyl)ethyl]-1-[(2R)-2-pyrrolidinylmethyl]-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide**



Following the general procedure in **Example 3** using intermediate 26 (100 mg, 0.200 mmol), 4-ethoxy- α -methyl-phenylacetyl chloride (46 mg, 0.230 mmol) and zinc dust (15 mg, 0.231 mmol) in 3 mL of 1,2-dichloroethane. The product was purified by reversed-phase HPLC using 10-50% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 25 mg (19%). ¹H NMR (CD₃OD) 1.32 (t, J = 7.03 Hz, 3H), 1.60 (m, 1H), 1.77 (d, J = 7.03 Hz, 3H), 1.96 (m, 2H), 2.11 (m, 1H), 3.10 (m, 1H), 3.24 (m, 1H), 3.37 (m, 1H), 3.73 (m, 1H), 3.98 (q, J = 7.03 Hz, 2H), 4.34 (m, 1H), 4.38 (m, 4H), 4.42 (m, 1H), 4.52 (m, 1H), 6.87 (d, J = 8.40 Hz, 2H), 7.15 (m, 2H), 7.41 (d, J = 8.40 Hz, 1H), 7.67 (d, J = 8.40 Hz, 1H), 7.81 (s, 1H); MS (ESI) 557.49 (MH⁺).

Example 22

2-[(5-Ethoxy-2-pyridyl)methyl]-1-[[2*R*]-1-methyl-2-piperidinyl]methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide




Intermediate 33 (135 mg, 0.263 mmol), HATU (120 mg, 0.315 mmol) and 5-ethoxy-2-pyridylacetic acid hydrochloride (70 mg, 0.315 mmol) were stirred in 5 mL of DMF containing DIPEA (0.095 mL, 0.526 mmol) at RT for 3h. The solution was diluted with EtOAc and washed with saturated NaHCO₃ solution, brine and dried over anhydrous Na₂SO₄. The solvent was evaporated. The residue was dissolved in 3 mL of 1,2-dichloroethane containing a catalytic amount of concentrated HCl (11.6 M) and the solution was stirred at 80 °C for 5h. The solution was cooled to RT and then

diluted with dichloromethane. The solution was washed with saturated NaHCO_3 solution, brine and dried over anhydrous Na_2SO_4 . The solvent was evaporated. The residue was dissolved in 5 mL of THF containing a catalytic amount of glacial acetic acid. An excess of 37% $\text{HCHO}/\text{H}_2\text{O}$ (1 mL) was added followed by $\text{NaBH}(\text{OAc})_3$ (68 mg, 0.316 mmol). The solution was stirred at RT for 1h. The solution was then diluted with EtOAc and washed with saturated NaHCO_3 solution, brine and dried over anhydrous Na_2SO_4 . The product was purified by reversed-phase HPLC using 10-50% $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ and lyophilized affording the title compound as the corresponding TFA salt. Yield: 53 mg (25%). ^1H NMR (CD_3OD) 1.39 (t, $J = 7.03$ Hz, 3H), 1.42 (m, 1H), 1.56 (m, 2H), 1.79 (m, 2H), 1.90 (m, 1H), 3.11 (brs, 4H), 3.60 (m, 1H), 3.80 (m, 1H), 4.12 (q, $J = 6.90$ Hz, 2H), 4.36 (brd, 4H), 4.57 (m, 1H), 4.63 (m, 1H), 4.86 (m, 1H), 5.10 (m, 1H), 7.44 (d, $J = 8.40$ Hz, 1H), 7.53 (s, 2H), 7.69 (s, 1H), 7.77 (d, $J = 8.40$ Hz, 1H), 8.22 (s, 1H); MS ESI: 572.25 (MH⁺).

Example 23

2-[(5-Ethoxy-2-pyridyl)methyl]-1-[[*(2R)*-1-methyl-2-pyrrolidinyl)methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide




Intermediate 26 (101 mg, 0.245 mmol), HATU (112 mg, 0.294 mmol) and 5-ethoxy-2-pyridylacetic acid hydrochloride (65 mg, 0.294 mmol) were stirred in DMF (5 mL) containing DIPEA (0.090 mL, 0.490 mmol) at RT for 3h. The solvent was evaporated. The residue was dissolved in EtOAc and washed with saturated NaHCO_3 solution, brine and dried over anhydrous Na_2SO_4 . The solvent was evaporated and the residue was dissolved in 1,2-dichloroethane (3 mL). A catalytic amount of concentrated HCl (11.6 M) was added and the solution was then stirred at 80 °C for 5h. The solution was cooled to RT and diluted with dichloromethane. The organic phase was washed with saturated NaHCO_3 , brine and dried over anhydrous Na_2SO_4 . The product was purified by reversed-phase HPLC using 10-50% $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ and

then lyophilized affording the title compound as the corresponding TFA salt. Yield: 86 mg (45%); ¹H NMR (CD₃OD) 1.39 (t, J = 6.93Hz, 3H), 1.98 (m, 1H), 2.15 (m, 2H), 2.29 (m, 1H), 2.86 (s, 3H), 3.32 (m, 1H), 3.85 (m, 1H), 4.12 (q, J = 6.90Hz, 2H), 4.19 (m, 1H), 4.36 (brs, 5H), 4.81 (m, 1H), 5.02 (m, 1H), 7.47 (d, J = 8.50Hz, 1H), 7.60 (m, 2H), 7.70 (s, 1H), 7.86 (d, J = 8.59Hz, 1H), 8.23 (s, 1H); MS (ESI) 558.19 (MH⁺).

Example 24


1-(Cyclobutylmethyl)-2-(4-ethoxybenzyl)-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide



Following the general procedure in Example 16 using intermediate 36 (113 mg, 0.295 mmol), 4-ethoxyphenylacetic acid (58 mg, 0.325 mmol), HATU (123 mg, 0.325 mmol) and DIPEA (0.075 mL, 0.443 mmol) in 5 mL of DMF. The final dehydrated product was purified by reversed-phase HPLC using 20-80% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 118 mg (62%); ¹H NMR (400 MHz, CD₃OD) δ 1.35 (t, J = 6.84 Hz, 3H), 1.86 (m, 4H), 1.97 (m, 2H), 2.77 (m, 1H), 4.01 (q, J = 7.03 Hz, 2H), 4.35 (m, 4H), 4.47 (d, J = 7.03 Hz, 2H), 4.52 (s, 2H), 6.93 (d, J = 8.59 Hz, 2H), 7.25 (d, J = 8.59 Hz, 2H), 7.56 (d, J = 8.59 Hz, 1H), 7.75 (s, 1H), 7.96 (d, J = 8.40 Hz, 1H); MS (ESI) 528.1 (MH⁺); Anal. Calcd for C₂₆H₂₇N₃O₂F₆ + 0.6TFA + 0.4H₂O: C, 54.17; H, 4.75; N, 6.97. Found: C, 54.08; H, 4.69; N, 6.96.

Example 25


1-(Cyclobutylmethyl)-2-[(5-ethoxypyridin-2-yl)methyl]-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide



Following the general procedure in **Example 16** using **intermediate 36** (119 mg,
 0.310 mmol), 5-ethoxy-2-pyridylacetic acid hydrochloride (75 mg, 0.341 mmol),
 HATU (130 mg, 0.341 mmol) and DIPEA (0.110 mL, 0.620 mmol) in 5 mL of DMF.
 The final dehydrated product was purified by reversed-phase HPLC using 20-80%
 CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding
 TFA salt. Yield: 135 mg (68%); ¹H NMR (400 MHz, CD₃OD) δ 1.40 (t, J = 7.03 Hz,
 3H), 1.87 (m, 4H), 2.00 (m, 2H), 2.83 (m, 1H), 4.11 (q, J = 6.96 Hz, 2H), 4.37 (brs,
 4H), 4.51 (d, J = 7.23 Hz, 2H), 4.72 (d, J = 7.23 Hz, 2H), 7.50 (m, 2H), 7.57 (d, J =
 8.59 Hz, 1H), 7.78 (s, J = 1H), 7.96 (d, J = 8.40 Hz, 1H), 8.21 (d, J = 2.54 Hz, 1H);
 MS (ESI) 529.1 (MH⁺); Anal. Calcd for C₂₅H₂₆N₄O₂F₆ + 1.5TFA + 0.2H₂O: C,
 47.83; H, 4.00; N, 7.97. Found: C, 47.80; H, 4.05; N, 7.93.

Example 26

1-(Cyclopentylmethyl)-2-[(5-ethoxypyridin-2-yl)methyl]-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide




Following the general procedure in **Example 16** using **intermediate 37** (129 mg,
 0.324 mmol), 5-ethoxy-2-pyridylacetic acid hydrochloride (78 mg, 0.356 mmol),
 HATU (135 mg, 0.356 mmol) and DIPEA (0.115 mL, 0.648 mmol) in 5 mL of DMF.
 The final dehydrated product was purified by reversed-phase HPLC using 20-80%

CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 155 mg (73%); ¹H NMR (400 MHz, CD₃OD) δ 1.30 (m, 2H), 1.40 (t, J = 6.83 Hz, 3H), 1.56 (m, 2H), 1.72 (m, 4H), 2.41 (m, 1H), 4.11 (q, J = 7.03 Hz, 2H), 4.38 (m, 4H), 4.43 (d, J = 7.81 Hz, 2H), 4.71 (s, 2H), 7.50 (s, 2H), 7.57 (d, J = 8.79 Hz, 1H), 7.77 (s, 1H), 7.95 (d, J = 8.59 Hz, 1H), 8.21 (s, 1H); MS (ESI) 543.1 (MH⁺); Anal. Calcd for C₂₆H₂₈N₄O₂F₆ + 1.1TFA: C, 50.71; H, 4.39; N, 8.39. Found: C, 50.76; H, 4.11; N, 8.36.

Example 27


2-(4-Ethoxybenzyl)-1-[(2*S*)-piperidin-2-ylmethyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide



Following the general procedure in Example 16 using intermediate 35 (138 mg, 0.269 mmol), 4-ethoxyphenylacetic acid (53 mg, 0.296 mmol), HATU (112 mg, 0.296 mmol) and DIPEA (0.050 mL, 0.404 mmol) in 5 mL of DMF. The final dehydrated product was purified by reversed-phase HPLC using 10-50% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 117 mg (65%); ¹H NMR (400 MHz, CD₃OD) δ 1.36 (t, J = 6.83 Hz, 3H), 1.44 (m, 1H), 1.57 (m, 1H), 1.70 (m, 1H), 1.86 (m, 3H), 2.79 (m, 1H), 3.37 (m, 2H), 4.01 (q, J = 7.03 Hz, 2H), 4.39 (m, 4H), 4.44 (s, 2H), 4.53 (m, 2H), 6.92 (d, J = 8.79 Hz, 2H), 7.22 (d, J = 8.79 Hz, 2H), 7.51 (d, J = 8.29 Hz, 1H), 7.76 (s, 1H), 7.80 (d, J = 8.40 Hz, 1H); MS (ESI) 557.1 (MH⁺).

Example 28

2-[(5-Ethoxypyridin-2-yl)methyl]-1-(3-furylmethyl)-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide




Following the general procedure in **Example 16** using 3-amino-4-[(3-furylmethyl)amino]-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (80 mg, 0.202 mmol), 5-ethoxy-2-pyridylacetic acid hydrochloride (48 mg, 0.222 mmol), HATU (85 mg, 0.222 mmol) and DIPEA (0.053 mL, 0.303 mmol) in 5 mL of DMF. The final dehydrated product was purified by reversed-phase HPLC using 20-80% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt.

Yield: 55 mg (42%); ¹H NMR (400 MHz, CD₃OD) δ 1.40 (t, J = 7.03 Hz, 3H), 4.11 (q, J = 7.03 Hz, 2H), 4.36 (m, 4H), 4.88 (s, 2H), 5.56 (s, 2H), 6.25 (s, 1H), 7.44 (s, 1H), 7.47 (s, 2H), 7.52 (m, 2H), 7.78 (s, 1H), 7.87 (d, J = 8.59 Hz, 1H), 8.20 (s, 1H); MS (ESI) 541.1 (MH⁺); Anal. Calcd for C₂₅H₂₂N₄O₃F₆ + 1.2TFA + 0.1 H₂O: C, 48.46; H, 3.47; N, 8.25. Found: C, 48.50; H, 3.44; N, 8.27.

Example 29

2-[(5-Ethoxypyridin-2-yl)methyl]-1-(3-thienylmethyl)-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide




Following the general procedure in **Example 16** using 3-amino-4-[(3-thienylmethyl)amino]-*N,N*-bis(2,2,2-trifluoroethyl)benzamide (125 mg, 0.303 mmol), 5-ethoxy-2-pyridylacetic acid hydrochloride (73 mg, 0.333 mmol), HATU (127 mg, 0.333 mmol) and DIPEA (0.105 mL, 0.606 mmol) in 5 mL of DMF. The final dehydrated product was purified by reversed-phase HPLC using 20-80% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt.

Yield: 137 mg (67%); ^1H NMR (400 MHz, CD_3OD) δ 1.38 (t, $J = 7.03$ Hz, 3H), 4.07 (q, $J = 7.03$ Hz, 2H), 4.36 (m, 4H), 4.90 (s, 2H), 5.65 (s, 2H), 6.86 (d, $J = 4.68$ Hz, 1H), 7.14 (s, 1H), 7.38 (m, 3H), 7.45 (d, $J = 8.59$ Hz, 1H), 7.73 (d, $J = 8.40$ Hz, 1H), 7.77 (s, 1H), 8.12 (s, 1H); MS (ESI) 557.0 (MH $^+$); Anal. Calcd for $\text{C}_{25}\text{H}_{22}\text{N}_4\text{O}_2\text{SF}_6 + 0.7 \text{ TFA} + 0.5 \text{ H}_2\text{O}$: C, 49.13; H, 3.70; N, 8.68. Found: C, 49.08; H, 3.75; N, 8.65.

Example 30

1-(Cyclohexylmethyl)-2-[(5-ethoxypyridin-2-yl)methyl]-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide



10


Following the general procedure in Example 16 using intermediate 24 (111 mg, 0.270 mmol), 5-ethoxy-2-pyridylacetic acid hydrochloride (65 mg, 0.297 mmol), HATU (113 mg, 0.297 mmol) and DIPEA (0.120 mL, 0.675 mmol) in 5 mL of DMF. The final dehydrated product was purified by reversed-phase HPLC using 20-80% $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 109 mg (60%); ^1H NMR (400 MHz, CD_3OD) δ 1.12 (m, 5H), 1.38 (t, $J = 6.93$ Hz, 3H), 1.59 (m, 2H), 1.60 (m, 1H), 1.68 (m, 2H), 1.78 (m, 1H), 4.10 (q, $J = 6.90$ Hz, 2H), 4.28 (d, $J = 7.62$ Hz, 2H), 4.36 (m, 4H), 4.70 (d, $J = 7.23$ Hz, 1H), 4.88 (m, 1H), 7.50 (m, 2H), 7.55 (dd, $J = 1.37, 8.59$ Hz, 1H), 7.77 (s, 1H), 7.93 (d, $J = 8.59$ Hz, 1H), 8.20 (d, $J = 2.73$ Hz, 1H); MS (ESI) 556.7 (MH $^+$); Anal. Calcd for $\text{C}_{27}\text{H}_{30}\text{N}_4\text{O}_2\text{F}_6 + 2.1 \text{ TFA} + 0.1 \text{ H}_2\text{O}$: C, 46.97; H, 4.08; N, 7.02. Found: C, 46.95; H, 4.12; N, 7.07.

20

Example 31

1-(Cyclohexylmethyl)-2-[(5-isopropoxypyridin-2-yl)methyl]-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide


25



Following the general procedure in **Example 16** using **intermediate 24** (140 mg, 0.340 mmol), 4-isopropoxy-2-pyridylacetic acid hydrochloride • 2LiCl (120 mg, 0.374 mmol), HATU (145 mg, 0.374 mmol) and DIPEA (0.175 mL, 1.02 mmol) in 5 mL of DMF. The final dehydrated product was purified by reversed-phase HPLC using 20-80% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 80 mg (35%); ¹H NMR (400 MHz, CD₃OD) δ 1.10 (m, 6H), 1.30 (d, J = 6.05 Hz, 6H), 1.57 (m, 2H), 1.62 (m, 1H), 1.70 (m, 2H), 1.74 (m, 1H), 4.23 (d, J = 7.62 Hz, 2H), 4.36 (m, 4H), 4.64 (m, 2H), 7.44 (s, 2H), 7.50 (dd, J = 1.37, 8.40 Hz, 1H), 7.73 (s, 1H), 7.84 (d, J = 8.59 Hz, 1H), 8.15 (s, 1H); MS (ESI) 571.2 (MH⁺); Anal. Calcd for C₂₈H₃₂N₄O₂F₆ + 0.6TFA + 0.2 H₂O: C, 54.58; H, 5.18; N, 8.72. Found: C, 54.55; H, 5.18; N, 8.67.

Example 32

2-(4-Ethoxybenzyl)-1-[(4-methylmorpholin-3-yl)methyl]-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide




Following the general procedure in **Example 22** using **intermediate 34** (135 mg, 0.262 mmol), 4-ethoxyphenylacetic acid (57 mg, 0.314 mmol), HATU (120 mg, 0.314 mmol) and DIPEA (0.091 mL, 0.524 mmol) in 5 mL of DMF. The reductive amination step was performed using NaBH(OAc)₃ (111 mg, 0.525 mmol) in 5 mL of THF. The final product was purified by reversed-phase HPLC using 10-50% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding

TFA salt. Yield: 103 mg (57%); ^1H NMR (400 MHz, CD_3OD) δ 1.34 (t, $J = 7.03$ Hz, 3H), 3.03 (s, 3H), 3.20 (m, 1H), 3.54 (m, 4H), 3.91 (m, 2H), 4.00 (q, $J = 7.03$ Hz, 2H), 4.35 (m, 4H), 4.40 (s, 2H), 4.72 (m, 2H), 6.92 (d, $J = 8.59$ Hz, 2H), 7.19 (d, $J = 8.79$ Hz, 2H), 7.45 (d, $J = 8.40$ Hz, 1H), 7.75 (m, 2H); MS (ESI) 573.2 (MH $^+$); Anal.

- 5 Calcd for $\text{C}_{27}\text{H}_{30}\text{N}_4\text{O}_3\text{F}_6 + 1.8$ TFA: C, 47.25; H, 4.12; N, 7.20. Found: C, 47.22; H, 4.00; N, 7.44.

Example 33


- 2-[(5-Ethoxypyridin-2-yl)methyl]-1-[(4-methylmorpholin-3-yl)methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide



- Following the general procedure in **Example 22** using **intermediate 34** (135 mg, 0.262 mmol), 5-ethoxy-2-pyridylacetic acid hydrochloride (68 mg, 0.314 mmol), HATU (120 mg, 0.314 mmol) and DIPEA (0.091 mL, 0.524 mmol) in 5 mL of DMF. The reductive amination step was performed by using $\text{NaBH}(\text{OAc})_3$ (111 mg, 0.525 mmol) in 5 mL of THF. The final product was purified by reversed-phase HPLC using 10-50% $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 46 mg (26%); ^1H NMR (400 MHz, CD_3OD) δ 1.38 (t, $J = 7.03$ Hz, 3H), 3.17 (s, 3H), 3.33 (m, 1H), 3.67 (m, 2H), 3.74 (m, 1H), 3.98 (m, 4H), 4.10 (q, $J = 7.03$ Hz, 2H), 4.35 (brs, 4H), 4.98 (m, 2H), 7.43 (d, $J = 8.40$ Hz, 1H), 7.51 (s, 2H), 7.68 (s, 1H), 7.78 (d, $J = 8.40$ Hz, 2H), 8.21 (s, 1H); MS (ESI) 574.2 (MH $^+$).

Example 34


2-(4-Ethoxybenzyl)-1-[(2*S*)-1-methylpiperidin-2-ylmethyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide



Following the general procedure in **Example 22** using **intermediate 35** (288 mg, 0.561 mmol), 4-ethoxyphenylacetic acid (120 mg, 0.673 mmol), HATU (255 mg, 0.673 mmol) and DIPEA (0.145 mL, 0.842 mmol) in 10 mL of DMF. The reductive amination step was performed by using NaBH(OAc)₃ (235 mg, 1.12 mmol) in 5 mL of THF. The final product was purified by reversed-phase HPLC using 10-50% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 257 mg (67%); ¹H NMR (400 MHz, CD₃OD) δ 1.21 (m, 1H), 1.34 (t, J = 6.93 Hz, 3H), 1.47 (m, 1H), 1.72 (m, 2H), 1.83 (m, 1H), 2.96 (m, 1H), 3.01 (s, 3H), 3.22 (m, 1H), 3.55 (m, 1H), 4.00 (q, J = 7.03 Hz, 2H), 4.37 (m, 5H), 4.40 (s, 2H), 4.86 (m, 1H), 6.92 (d, J = 8.59 Hz, 2H), 7.20 (d, J = 8.59 Hz, 2H), 7.45 (d, J = 8.40 Hz, 1H), 7.72 (m, 1H), 7.75 (s, 1H); MS (ESI) 571.2 (MH⁺); Anal. Calcd for C₂₈H₃₂N₄O₂F₆ + 1.8 TFA + 0.3 H₂O: C, 47.07; H, 4.24; N, 6.78. Found: C, 47.03; H, 4.20; N, 6.93.

Example 35

2-(4-Isopropoxybenzyl)-1-{[(2*R*)-1-methylpiperidin-2-yl]methyl}-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide




Following the general procedure in **Example 22** using **intermediate 33** (124 mg, 0.242 mmol), 4-isopropoxyphenylacetic acid (52 mg, 0.266 mmol), HATU (102 mg, 0.266 mmol) and DIPEA (0.065 mL, 0.363 mmol) in 5 mL of DMF. The reductive amination step was performed by using NaBH(OAc)₃ (105 mg, 0.484 mmol) in 5 mL

of THF. The final product was purified by reversed-phase HPLC using 10-50% CH₃CN/H₂O and then lyophilized affording the title compound as the corresponding TFA salt. Yield: 112 mg (66%); ¹H NMR (400 MHz, CD₃OD) δ 1.19 (m, 1H), 1.26 (d, J = 6.05 Hz, 6H), 1.34 (m, 1H), 1.45 (m, 1H), 1.70 (m, 2H), 1.81 (m, 1H), 2.96 (m, 1H), 3.01 (s, 3H), 3.56 (m, 1H), 4.36 (m, 5H), 4.40 (s, 2H), 4.56 (dt, J = 6.05 Hz, 1H), 4.88 (m, 1H), 6.90 (d, J = 8.79 Hz, 2H), 7.20 (d, J = 8.79 Hz, 2H), 7.45 (dd, J = 1.37, 8.59 Hz, 1H), 7.74 (m, 2H); MS (ESI) 585.2 (MH⁺); Anal. Calcd for C₂₉H₃₄N₄O₂F₆ + 2.3 TFA + 0.1 H₂O: C, 47.55; H, 4.34; N, 6.60. Found: C, 47.51; H, 4.33; N, 6.74.

What is claimed is:

1. A compound of formula (I) or pharmaceutically acceptable salts thereof:



wherein

- 5 R^{F1} and R^{F2} are independently electron-withdrawing groups;

Z is selected from O= and S=;

- R^1 is selected from C_{1-10} alkyl; C_{1-10} alkyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; C_{2-10} alkenyl; C_{2-10} alkenyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; C_{2-10} alkynyl; C_{2-10} alkynyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; R^3R^4N - C_{1-6} alkyl; $R^3R^4NC(=O)-C_{1-6}$ alkyl; R^3O-C_{1-6} alkyl; $R^3OC(=O)-C_{1-6}$ alkyl; $R^3C(=O)-C_{1-6}$ alkyl; $R^3C(=O)NR^3-C_{1-6}$ alkyl; $R^3R^4NSO_2-C_{1-6}$ alkyl; $R^3CSO_2N(R^4)-C_{1-6}$ alkyl; $R^3R^4NC(=O)N(R^5)-C_{1-6}$ alkyl; $R^3R^4NSO_2N(R^5)-C_{1-6}$ alkyl; aryl- C_{1-6} alkyl; aryl- $C(=O)-C_{1-6}$ alkyl; heterocyclyl- C_{1-6} alkyl; heterocyclyl- $C(=O)-C_{1-6}$ alkyl; substituted aryl- C_{1-6} alkyl; substituted aryl- $C(=O)-C_{1-6}$ alkyl; substituted heterocyclyl- C_{1-6} alkyl; substituted heterocyclyl- $C(=O)-C_{1-6}$ alkyl; and C_{1-10} hydrocarbylamino;

- R^2 is selected from C_{1-6} alkyl, substituted C_{1-6} alkyl, C_{2-6} alkenyl, substituted C_{2-6} alkenyl, C_{2-6} alkynyl, substituted C_{2-6} alkynyl, C_{3-6} cycloalkyl, substituted C_{3-6} cycloalkyl, aryl, substituted aryl, and C_{5-6} heteroaryl, and substituted C_{5-6} heteroaryl;

R^3 , R^4 and R^5 are independently selected from -H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and a divalent C_{1-6} group that together with another divalent C_{1-6} group forms a portion of a ring;

- X is a C_{1-10} divalent group that separates groups connected thereto by one or two atoms;

Ar is a C_{4-12} divalent aromatic group; and

Y is selected from -CH= and -N=.

2. The compound as claimed in claim 1, wherein

R^{F1} and R^{F2} are independently C_{1-6} alkyl substituted by one or more groups selected from -F, -Cl, -Br, -NO₂, -CN, -OH, -CHO, -C(=O)-R' and -OR', wherein R' is a C_{1-3} alkyl.

5 3. The compound as claimed in claim 1, wherein

R^{F1} and R^{F2} are independently selected from -CF₃, -CH₂CF₃, -CH₂CHF₂, -CHFCH₂F, -CHFCHF₂, -CHFCH₂F, -CF₂CF₃, -CF₂CH₃, -CF₂CH₂F, -CF₂CHF₂, -CF₃, -CH₂CCl₃, -CH₂CHCl₂, -CH₂CBr₃, -CH₂CHBr₂, -CH₂NO₂, -CH₂CH₂NO₂, -CH₂CN, -CH₂CH₂CN, and -CH₂CH₂OCH₃.

10

4. The compound as claimed in claim 1, wherein R^{F1} and R^{F2} are independently C_{1-6} groups that comprise at least 30% fluorine by weight and Z is O=.

5. The compound as claimed in claim 1, wherein R^1 is selected from C_{1-10} alkyl; C_{1-10} alkyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; C_{2-10} alkenyl; C_{2-10} alkenyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; C_{2-10} alkynyl; C_{2-10} alkynyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; $R^3R^4N-C_{1-6}$ alkyl; $R^3R^4NC(=O)-C_{1-6}$ alkyl; R^3O-C_{1-6} alkyl; $R^3OC(=O)-C_{1-6}$ alkyl; $R^3C(=O)-C_{1-6}$ alkyl; $R^3C(=O)NR^3-C_{1-6}$ alkyl; $R^3R^4NSO_2-C_{1-6}$ alkyl; $R^3CSO_2N(R^4)-C_{1-6}$ alkyl; $R^3R^4NC(=O)N(R^5)-C_{1-6}$ alkyl; $R^3R^4NSO_2N(R^5)-C_{1-6}$ alkyl; aryl- C_{1-6} alkyl; aryl-C(=O)- C_{1-6} alkyl; heterocyclyl- C_{1-6} alkyl; heterocyclyl-C(=O)- C_{1-6} alkyl; substituted aryl- C_{1-6} alkyl; substituted aryl-C(=O)- C_{1-6} alkyl; substituted heterocyclyl- C_{1-6} alkyl; substituted heterocyclyl-C(=O)- C_{1-6} alkyl; and C_{1-10} hydrocarbylamino;

25 R^2 is selected from C_{1-6} alkyl, C_{1-6} alkyl substituted by at least one fluorine, C_{2-6} alkenyl, C_{2-6} alkenyl substituted by at least one fluorine, C_{2-6} alkynyl, C_{2-6} alkynyl substituted by at least one fluorine, C_{3-6} cycloalkyl, substituted C_{3-6} cycloalkyl, aryl, substituted aryl, and C_{5-6} heteroaryl, and substituted C_{5-6} heteroaryl;

30 R^3 , R^4 and R^5 are independently selected from -H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and a divalent C_{1-6} group that together with another divalent C_{1-6} group forms a portion of a ring; and

X is selected from $-\text{NR}^6-$, $-\text{C}(=\text{O})-$, $-\text{CH}_2-\text{CH}_2-$, $-\text{CH}=\text{CH}-$, $-\text{O}-$, $-\text{C}(\text{R}^6)(\text{R}^7)-$, and $-\text{S}(\text{O})_n-$, wherein n is 0, 1 or 2, wherein R^6 and R^7 are independently C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, $-\text{OH}$, or $-\text{H}$.

- 5 6. A compound according to Claim 1,
wherein:

R^1 is selected from C_{1-8} alkyl; C_{2-8} alkenyl; C_{2-8} alkynyl; aryl- C_{1-6} alkyl; aryl- C_{1-6} alkyl with the aryl substituted by at least one group selected from C_{1-6} alkyl, acetoxymethyl, nitro and halogen; $\text{R}^8\text{R}^9\text{NC}_{1-6}$ alkyl; $\text{R}^8\text{OC}_{1-6}$ alkyl; cycloalkyl-
10 C_{1-6} alkyl; heterocycloalkyl- C_{1-6} alkyl; heterocycloalkyl- C_{1-6} alkyl with the heterocycloalkyl thereof substituted by at least one group selected from C_{1-8} alkyl, acetoxymethyl, nitro and halogen; C_{1-6} alkylaryl; C_{1-6} alkyl- $\text{C}(=\text{O})-$; C_{6-8} aryl- $\text{C}(=\text{O})-$; C_{4-8} heteroaryl- $\text{C}(=\text{O})-$; heteroaryl- C_{1-6} alkyl; heteroaryl- C_{1-6} alkyl with the heteroaryl thereof substituted by at least one group selected from C_{1-6} alkyl, acetoxymethyl, nitro
15 and halogen; and R^NC_{1-6} alkyl;

R^2 is selected from $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)_2$, C_{3-6} cycloalkyl, $-\text{CH}_2\text{CF}_3$, $-\text{CHF}_2$, $-\text{CF}_3$ and aryl;

R^N is an oxidized pyridyl wherein the nitrogen atom on the pyridyl ring is in an oxidized state (N^+-O^-);

20 Ar is selected from an arylene; an heteroarylene; an arylene substituted by at least one group selected from C_{1-6} alkyl, halogen, trifluoromethyl, cyano, nitro, hydroxy and C_{1-6} alkoxy; and an heteroarylene substituted by at least one group selected from C_{1-6} alkyl, halogen, trifluoromethyl, cyano, nitro, hydroxy and C_{1-6} alkoxy; and

25 R^8 and R^9 are independently selected from $-\text{H}$ and C_{1-6} alkyl.

7. The compound according to claim 6,

wherein the arylene is *para*-arylene; and the heteroarylene is selected from six-membered ring *para*-heteroarylene and five-membered ring *meta*-heteroarylene.

30

8. A compound according to Claim 1,
wherein:

R^1 is selected from ethyl, propyl, allyl, isopentyl, benzyl, dimethylaminoethyl, 4-pyridylmethyl, 2-pyridylmethyl, 1-pyrrolylethyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, 2-pyrrolidylmethyl, 3-pyrrolidylmethyl, N-methyl-2-pyrrolidylmethyl, N-methyl-3-pyrrolidylmethyl, 2-piperidylmethyl, 3-piperidylmethyl, 4-piperidylmethyl, N-methyl-2-piperidylmethyl, N-methyl-3-piperidylmethyl, N-methyl-4-piperidylmethyl, 3-thienylmethyl, 2-tetrahydrofuranylmethyl, 3-tetrahydrofuranylmethyl, 2-tetrahydropyranylmethyl, 3-tetrahydropyranylmethyl, 4-tetrahydropyranylmethyl, (2-nitrothiophene-5-yl)methyl, (1-methyl-1H-imidazole-2-yl)methyl, (5-(acetoxymethyl)-2-furanylmethyl, (2,3-dihydro-1H-isoin-1-yl)methyl, and 5-(2-methylthiazolyl);

R^2 is selected from $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CF}_3$, CF_3 , cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and phenyl;

R^{F1} and R^{F2} are $-\text{CH}_2\text{CF}_3$ and Z is $\text{O}=\text{}$;

Ar is selected from a *para*-arylene; a *para*-arylene substituted with C_{1-6} alkyl, halogen, trifluoromethyl, cyano, nitro, hydroxy and C_{1-6} alkoxy; a six-membered ring *para*-heteroarylene; and a six-membered ring *para*-heteroarylene substituted with a group selected from C_{1-6} alkyl, halogen, trifluoromethyl, cyano, nitro, hydroxy and C_{1-6} alkoxy.

9. A compound according to Claim 1, wherein:

R^{F1} and R^{F2} are $-\text{CH}_2\text{CF}_3$, and Z is $\text{O}=\text{}$;

R^2 is $-\text{CH}_2\text{CH}_3$;

Ar is selected from *para*-phenylene and *para*-pyridylene; and

X is selected from $-\text{CH}_2-$ and $-\text{CH}(\text{CH}_3)-$.

10. A compound according to claim 1, wherein said compound is selected from:

2-[(4-Ethoxyphenyl)methyl]-1-(3-methylbutyl)-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide;

1-(Cyclopropylmethyl)-2-[(4-ethoxyphenyl)methyl]-N,N-bis(2,2,2-trifluoroethyl)-1H-benzimidazole-5-carboxamide;

- 1-(Cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 2-[(4-Ethoxyphenyl)methyl]-1-(2-furanylmethyl)-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 5 2-[(4-Ethoxyphenyl)methyl]-1-[(2*S*)-2-pyrrolidinylmethyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 2-[(4-Ethoxyphenyl)methyl]-1-[(2*R*)-2-pyrrolidinylmethyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 10 2-[(4-ethoxyphenyl)methyl]-1-(4-pyridinylmethyl)-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 2-[1-(4-Ethoxyphenyl)ethyl]-1-(4-pyridinylmethyl)-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 2-[(4-Ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 15 2-[(4-Ethoxyphenyl)methyl]-1-[(2*R*)-tetrahydro-2-furanyl]methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 2-[(4-Ethoxyphenyl)methyl]-1-[(2*S*)-tetrahydro-2-furanyl]methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 20 2-[(4-Ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-2-yl)methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 2-[(4-Ethoxyphenyl)methyl]-1-[(2*R*)-2-piperidinylmethyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 2-[(5-Ethoxy-2-pyridyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 25 2-[(5-Ethoxy-2-pyridinyl)methyl]-1-(3-methylbutyl)-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;

- 2-[(4-Ethoxyphenyl)methyl]-1-[(2*R*)-1-methyl-2-pyrrolidinyl]methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 2-[(4-Ethoxyphenyl)methyl]-1-[(2*R*)-1-methyl-2-piperidinyl]methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 5 2-[(5-Ethoxy-2-pyridinyl)methyl]-1-[(2*R*)-2-pyrrolidinylmethyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 2-[1-(4-Ethoxyphenyl)ethyl]-1-[(2*R*)-2-pyrrolidinylmethyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 10 2-[(5-Ethoxy-2-pyridinyl)methyl]-1-[(2*R*)-1-methyl-2-piperidinyl]methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 2-[(5-Ethoxy-2-pyridinyl)methyl]-1-[(2*R*)-1-methyl-2-pyrrolidinyl]methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 1-(Cyclobutylmethyl)-2-(4-ethoxybenzyl)-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 15 1-(Cyclobutylmethyl)-2-[(5-ethoxypyridin-2-yl)methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 1-(Cyclopentylmethyl)-2-[(5-ethoxypyridin-2-yl)methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 20 2-(4-Ethoxybenzyl)-1-[(2*S*)-piperidin-2-ylmethyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 2-[(5-Ethoxypyridin-2-yl)methyl]-1-(3-furylmethyl)-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 2-[(5-Ethoxypyridin-2-yl)methyl]-1-(3-thienylmethyl)-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;
- 25 1-(Cyclohexylmethyl)-2-[(5-ethoxypyridin-2-yl)methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;

1-(Cyclohexylmethyl)-2-[(5-isopropoxy-pyridin-2-yl)methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;

2-(4-Ethoxybenzyl)-1-[(4-methylmorpholin-3-yl)methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;

5 2-[(5-Ethoxy-pyridin-2-yl)methyl]-1-[(4-methylmorpholin-3-yl)methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;

2-(4-Ethoxybenzyl)-1-1-[(2*S*)-1-methylpiperidin-2-yl)methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;

10 2-(4-Isopropoxybenzyl)-1-1-[(2*R*)-1-methylpiperidin-2-yl)methyl]-*N,N*-bis(2,2,2-trifluoroethyl)-1*H*-benzimidazole-5-carboxamide;

and pharmaceutically acceptable salts thereof.

11. A compound according to any one of above claims for use as a medicament.

12. The use of a compound according to any one of claims 1-10 in the
15 manufacture of a medicament for the therapy of pain.


13. The use of a compound according to any one of claims 1-10 in the
manufacture of a medicament for the treatment of cancers.

20 14. The use of a compound according to any one of claims 1-10 in the
manufacture of a medicament for the treatment of multiple sclerosis, Parkinson's
disease, Huntington's chorea, transplant rejection or Alzheimer's disease.

15. A pharmaceutical composition comprising a compound according to any one
25 of claims 1-10 and a pharmaceutically acceptable carrier.

16. A method for the therapy of pain in a warm-blooded animal, comprising the
step of administering to said animal in need of such therapy a therapeutically effective
amount of a compound according to any one of claims 1-10.


17. A method of producing a compound comprising the step of reacting a compound represented by formula (II) with $R^2OArXCOA$:



wherein

- 5 R^{F1} and R^{F2} are independently electron-withdrawing groups;
 Z is selected from $O=$ and $S=$;
 R^1 is selected from C_{1-10} alkyl; C_{1-10} alkyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; C_{2-10} alkenyl; C_{2-10} alkenyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; C_{2-10} alkynyl; C_{2-10} alkynyl substituted by at least one of halogen, cyano, acetoxymethyl and nitro; R^3R^4N - C_{1-6} alkyl; $R^3R^4NC(=O)-C_{1-6}$ alkyl; R^3O-C_{1-6} alkyl; $R^3OC(=O)-C_{1-6}$ alkyl; $R^3C(=O)-C_{1-6}$ alkyl; $R^3C(=O)NR^3-C_{1-6}$ alkyl; $R^3R^4NSO_2-C_{1-6}$ alkyl; $R^3CSO_2N(R^4)-C_{1-6}$ alkyl; $R^3R^4NC(=O)N(R^5)-C_{1-6}$ alkyl; $R^3R^4NSO_2N(R^5)-C_{1-6}$ alkyl; aryl- C_{1-6} alkyl; aryl- $C(=O)-C_{1-6}$ alkyl; heterocyclyl- C_{1-6} alkyl; heterocyclyl- $C(=O)-C_{1-6}$ alkyl; substituted aryl- C_{1-6} alkyl; substituted aryl- $C(=O)-C_{1-6}$ alkyl; substituted heterocyclyl- C_{1-6} alkyl; substituted heterocyclyl- $C(=O)-C_{1-6}$ alkyl; and C_{1-10} hydrocarbylamino;
- 10 R^2 is selected from C_{1-6} alkyl, substituted C_{1-6} alkyl, C_{2-6} alkenyl, substituted C_{2-6} alkenyl, C_{2-6} alkynyl, substituted C_{2-6} alkynyl, C_{3-6} cycloalkyl, substituted C_{3-6} cycloalkyl, aryl, substituted aryl, and C_{5-6} heteroaryl, and substituted
- 15 C_{5-6} heteroaryl;
- R^3 , R^4 and R^5 are independently selected from -H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and a divalent C_{1-6} group that together with another divalent C_{1-6} group forms a portion of a ring;
- 20 X is a C_{1-10} divalent group that separates groups connected thereto by one or
- 25 two atoms;
- A is selected from -OH, -Cl, -Br, and -I;
- Ar is a C_{4-12} divalent aromatic group; and
- Y is selected from -CH= and -N=.

18. A method of producing a compound comprising the step of reacting a compound represented by formula (III) with formaldehyde:



wherein

- 5 r and s are selected from 0, 1 and 2;
 R¹⁰ is selected from C₁₋₆alkylene, -O-, and -NR¹¹-, wherein R¹¹ is a C₁₋₆alkyl;
 R^{F1} and R^{F2} are independently electron-withdrawing groups;
 X is a C₁₋₁₀divalent group that separates groups connected thereto by one or
 two atoms;
- 10 Ar is a C₄₋₁₂divalent aromatic group;
 R² is selected from C₁₋₆alkyl, substituted C₁₋₆alkyl, C₂₋₆alkenyl, substituted
 C₂₋₆alkenyl, C₂₋₆alkynyl, substituted C₂₋₆alkynyl, C₃₋₆cycloalkyl, substituted
 C₃₋₆cycloalkyl, aryl, substituted aryl, and C₅₋₆heteroaryl, and substituted
 C₅₋₆heteroaryl; and
- 15 Y is selected from -CH= and -N=.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2003/001604

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07D 235/06, C07D 235/12, C07D 401/06, C07D 401/14, C07D 403/06,
C07D 405/06, C07D 405/14, A61K 31/4184 31/4439, A61P 25/04, 25/16, 35/00
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D, A61K, A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FR 1481049 A (SERGE LÉCOLIER), 19 May 1967 (19.05.1967)	1-18
P,A	WO 02085866 A1 (ASTRAZENECA AB), 31 October 2002 (31.10.2002)	1-18
A	WO 0236590 A1 (AMRAD OPERATIONS PTY LTD), 10 May 2002 (10.05.2002)	1-18
A	WO 02060447 A1 (UNIVERSITY OF CONNECTICUT), 8 August 2002 (08.08.2002)	1-18

☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier application or patent but published on or after the international filing date	"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 26 January 2004	Date of mailing of the international search report 27-01-2004
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86	Authorized officer EVA JOHANSSON/BS Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2003/001604

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0445781 A1 (STERLING DRUG INC), 11 Sept 1991 (11.09.1991) --	1-18
A	WO 0046209 A1 (SANOFI-SYNTHELABO), 10 August 2002 (10.08.2002) --	1-18
A	WO 9719063 A1 (SANOFI), 29 May 1997 (29.05.1997) --	1-18
A	STN International, file CAPLUS, CAPLUS accession no. 1973:87, document no. 78:87, Paglietti G. et al, "Dialkylaminoalkylbenzimidazoles of pharmacological interest", & Studi Sassaressi, Sezione 2: Archivio Bimestrale di Scienze Mediche e Naturali (1971), 49(5-6), 192-203 -- -----	1-18

INTERNATIONAL SEARCH REPORT

International application No.
PCT SE2003/01604

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 16
because they relate to subject matter not required to be searched by this Authority, namely:
see extra sheet
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT SE2003/01604

Box II.1

Claim 16 relate to methods of treatment of the human or animal body by surgery or by therapy or diagnostic methods practiced on the human or animal body (Rule 39.1(iv)). Nevertheless, a search has been executed for this claim. The search has been based on the alleged effects of the compounds or compositions

INTERNATIONAL SEARCH REPORT
Information on patent family members

24/12/2003

International application No.

PCT/SE 2003/001604

FR	1481049	A	19/05/1967	NONE		
WO	02085866	A1	31/10/2002	AU	7477201 A	14/01/2002
				CA	2414995 A	10/01/2002
				EP	1307481 A	07/05/2003
WO	0236590	A1	10/05/2002	AU	1366802 A	15/05/2002
				AU	PR118000 D	00/00/0000
				CA	2427394 A	10/05/2002
				EP	1339710 A	03/09/2003
WO	02060447	A1	08/08/2002	CA	2436133 A	08/08/2002
				EP	1363632 A	26/11/2003
EP	0445781	A1	11/09/1991	SE	0445781 T3	
				AT	133663 T	15/02/1996
				AU	643757 B	25/11/1993
				AU	7266991 A	12/09/1991
				CA	2036307 A,C	09/09/1991
				DE	69116705 D,T	26/09/1996
				DK	445781 T	19/02/1996
				ES	2082872 T	01/04/1996
				FI	95369 B,C	13/10/1995
				FI	911131 A	09/09/1991
				GR	3019058 T	31/05/1996
				HU	56492 A	30/09/1991
				HU	215125 B	28/09/1998
				IE	72953 B	07/05/1997
				IE	910700 A	11/09/1991
				IL	97426 A	31/03/1996
				JP	3157179 B	16/04/2001
				JP	4235164 A	24/08/1992
				KR	183396 B	01/05/1999
				MX	24839 A	01/10/1993
				MX	174433 B	16/05/1994
				NO	179612 B,C	05/08/1996
				NO	910868 A	09/09/1991
				NZ	237309 A	25/02/1993
				PH	27022 A	01/02/1993
				PT	96938 A,B	31/10/1991
				RU	2073670 C	20/02/1997
				US	5013837 A	07/05/1991

INTERNATIONAL SEARCH REPORT

Information on patent family members

24/12/2003

International application No.

PCT/SE 2003/001604

WO	0046209	A1	10/08/2002	AT	214598	T	15/04/2002
				AT	252562	T	15/11/2003
				AU	754771	B	21/11/2002
				AU	758299	B	20/03/2003
				AU	2298900	A	25/08/2000
				AU	3830299	A	06/12/1999
				BG	104963	A	31/07/2001
				BG	105749	A	28/02/2002
				BR	0007895	A	30/10/2001
				BR	9910606	A	02/10/2001
				CA	2333188	A	25/11/1999
				CA	2358885	A	10/08/2000
				CN	1346349	T	24/04/2002
				CZ	20012697	A	17/10/2001
				DE	60006070	D	00/00/0000
				DE	69901062	D,T	31/10/2002
				DK	1077687	T	17/06/2002
				EE	200100399	A	15/10/2002
				EP	1077687	A,B	28/02/2001
				SE	1077687	T3	
				EP	1150961	A,B	07/11/2001
				FR	2789078	A,B	04/08/2000
				FR	2789079	A,B	04/08/2000
				HK	1033271	A	00/00/0000
				HR	20010564	A	31/08/2002
				HU	0102027	A	28/03/2002
				HU	0201278	A	28/12/2002
				IL	139774	D	00/00/0000
				IL	144198	D	00/00/0000
				JP	2002515423	T	28/05/2002
				JP	2002536366	T	29/10/2002
				NO	20005838	A	10/01/2001
				NO	20013736	A	28/09/2001
				NZ	508392	A	27/09/2002
				NZ	512886	A	25/10/2002
				PL	344314	A	22/10/2001
				PL	350030	A	21/10/2002
				SK	10872001	A	03/12/2001
				TR	200102054	T	00/00/0000
				US	6432984	B	13/08/2002
				US	6645985	B	11/11/2003
				US	2002188007	A	12/12/2002
				ZA	200105739	A	24/01/2002

INTERNATIONAL SEARCH REPORT

Information on patent family members

24/12/2003

International application No.

PCT/SE 2003/001604

WO	9719063	A1	29/05/1997	AT	203015	T	15/07/2001
				AU	7698596	A	11/06/1997
				BR	9611618	A	30/03/1999
				CA	2238430	A	29/05/1997
				DE	69613860	D,T	04/04/2002
				DK	876350	T	05/11/2001
				EP	0876350	A,B	11/11/1998
				SE	0876350	T3	
				ES	2161382	T	01/12/2001
				FR	2741621	A,B	30/05/1997
				GR	3036830	T	31/01/2002
				JP	2000500479	T	18/01/2000
				NO	310508	B	16/07/2001
				NO	982330	A	22/05/1998
				PT	876350	T	28/12/2001
				US	6028084	A	22/02/2000
